

Research on Poverty Alleviation, REPOA, is an independent, non-profit organization concerned with poverty and related policy issues in Tanzania. REPOA undertakes and facilitates research, enables monitoring, and promotes capacity building, dialogue and knowledge sharing.

REPOA's research agenda is concerned with poverty and its alleviation. Our objectives are to:

- develop the research capacity in Tanzania;
- enhance stakeholders' knowledge of poverty issues and empower them to act;
- contribute to policy dialogue;
- support the monitoring of the implementation of poverty related policy;
- strengthen national and international poverty research networks, and forge linkages between research(ers) and users.

It is our conviction that research provides the means for the acquisition of knowledge necessary for improving the quality of welfare in Tanzanian society.

REPOA's Research Reports contain the result of research financed by REPOA. Our Special Papers contain the findings of commissioned studies conducted under our programmers of research, training and capacity building. The authors of these research reports and special papers are entitled to use their material in other publications; with acknowledgement to REPOA.

REPOA has published the results from this research as part of our mandate to disseminate information. Any views expressed are those of the authors alone and should not be attributed to REPOA.

Research on Poverty Alleviation (REPOA)
P.O. Box 33223, Dar es Salaam, Tanzania
157 Mgombani Street, Regent Estate
Tel: +255(0)(22) 270 00 83 / 277 25 56
Fax: +255(0)(22) 277 57 38
Email: repa@repa.or.tz
Website: www.repa.or.tz

ISBN: 978 - 9987 - 615 - 45 - 2



Research Report 10/1

Coping Strategies Used by Street Children in the Event of Illness

By Zena Amury
and Aneth Komba

RESEARCH ON POVERTY
ALLEVIATION

Coping Strategies Used by Street Children in the Event of Illness

By Zena Amury and Aneth Komba

Research Report 10/1



Published for: Research on Poverty Alleviation (REPOA)
P.O. Box 33223, Dar es Salaam, Tanzania
157 Mgombani Street, Regent Estate
Tel: +255(0) (22) 2700083 / 2772556
Fax: +255(0) (22) 2775738
Email: repa@repa.or.tz
Website: www.repa.or.tz

Design: Total Identity Ltd.

Suggested Citation:
Zena Amury and Aneth Komba
'Coping Strategies Used by Street Children in the Event of Illness'
Research Report 10/1, Dar es Salaam, REPOA

Suggested Keywords: Street children, illness, diseases, coping strategies, healthcare

© REPOA, 2010

ISBN: 978-9987-615-45-2

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means without the written permission of the copyright holder or the publisher.

Table of Contents

List of Table	v
List of Figures	vi
Abbreviations	vii
Acknowledgements	iv
Abstract	ix
1.0 Introduction	1
1.1 Background: The phenomenon of street children	1
1.2 Research problem and significance of the study	2
1.3 Research objectives	3
1.4 Research questions	3
2.0 Literature Review	4
2.1 Definition of children living on the streets	4
2.2 Tanzania's National Health Policy and its implications for access to healthcare	4
2.3 Accessibility of healthcare and health-seeking behaviour of children living on the streets	5
2.4 Vulnerability of street children to diseases and illnesses	6
2.5 The gender dimension of vulnerability	7
3.0 Methodology	8
3.1 Research approach	8
3.2 Study locations	8
3.3 Study participants and sampling procedure	9
3.4 Survey instruments and data collection	9
3.5 Data analysis	11
3.6 Ethical considerations	11
3.7 Limitations of the study	11
4. Findings and Discussions	13
4.1 General characteristics of respondents	13
4.2 Common diseases and illnesses of street children	14
4.3 Health-seeking behaviour/coping strategies used by street children	16
4.4 Children's reasons for not using health services	20
4.5 Children's suggestions to improve healthcare access	21
4.6 Sexual behaviours that put street children at risk of contracting HIV/AIDS and other sexually transmitted diseases	22
4.7 Activities that put street children at risk of contracting diseases and illnesses	23

Table of Contents

5.0	Summary of Key Findings, Conclusions and Recommendations	25
5.1	Summary of key findings	25
5.2	Conclusions	25
5.3	Policy recommendations	26
5.4	Recommendations for practical action to support street children	26
5.5	Recommendations for future research	28
	Bibliography and References	29
	Appendices	35
	Appendix 1: Findings from the control sample	35
	Appendix 2: Swahili questionnaire	35
	Appendix 3: English questionnaire	39
	Appendix 4: Semi-structured interview guide	42
	Appendix 5: Participatory exercise guide	44
	Publications by REPOA	46

Lists of Tables

Table 1	Number of participants	9
Table 2	General characteristics of respondents	13
Table 3	Children who experienced sickness during the last three months	15
Table 4	Diseases and illnesses frequently experienced by street children	15
Table 5	Health-seeking behaviour/coping strategies used by street children when they get sick	16
Table 6	Children's reasons for going to the hospital when sick	19
Table 7	Children's reasons for buying medicine when sick	19
Table 8	Children's reasons for sleeping off their illnesses	19
Table 9	Children's reasons for not going to the hospital	21
Table 10	Children's suggestions to improve healthcare access	22
Table 11	Children's sexual behaviour	22
Table 12	Income-earning activities of street children	24
Table A1	General characteristics of the control sample	35
Table A2	Key findings from the control sample	35

List of Figures

Figure 1	Spider diagram showing the types of diseases/illnesses suffered by participants	16
Figure 2	Drawing showing a boy going to the pharmacy	17
Figure 3	Drawing showing a boy sleeping when he was sick	18
Figure 4	Drawing showing a girl being chased away from the hospital	21

Abbreviations

CBO	Community-Based Organisation
ILO	International Labour Organisation
MCDGC	Ministry of Community Development, Gender and Children
MCH	Maternal and Child Health
MOH	Ministry of Health
NGO	Non-Governmental Organisation
OVC	Orphans and Vulnerable Children
PA	Participatory Approach
SAP	Structural Adjustment Policies
SP	Sulfadoxine-Pyrimethamine
SPSS	Statistical Package for Social Sciences
SRH	Sexual and Reproductive Health
STDs	Sexually Transmitted Diseases
STIs	Sexually Transmitted Infections

Acknowledgements

We are very grateful to Research on Poverty Alleviation (REPOA) for funding this research study, and for their constructive and tireless guidance throughout the project.

We would also like to extend our appreciation to the Directors of Ilala Municipal Council, Kinondoni Municipal Council and Temeke Municipal Council for allowing the research to be conducted in their municipalities.

We could not have completed the study without the support of research assistants who helped with data collection and analysis. Their support and contribution are greatly appreciated.

Finally, this study would have not been possible without the involvement of the street children who agreed to share their experiences. God bless them all.

***‘To the children living on the streets...who are struggling
to become what they should have been’***

Zena Amury and Aneth Komba

Abstract

Worldwide, street children are highly vulnerable to illness, disease and injury but tend to under-utilise health services. It is therefore critical to understand the coping strategies street children use when ill and the barriers they face in accessing health services.

This study examined the health-seeking behaviours of street children in three municipal districts of Dar es Salaam, Tanzania. The research was carried out in October 2005 and involved 272 street children (163 boys and 109 girls) aged 10 to 18 years. Both quantitative and qualitative methodologies were used for data collection.

Findings reveal that street children frequently experience illnesses and injuries, including fever, skin diseases, headaches, respiratory infections, diarrhoea and stomach upsets. However, the majority of children did not use health services. The cost of services and unfriendly attitudes of health workers were the barriers to access most often cited by participants. The children preferred to buy medicine from local shops and pharmacies because it is cheap and saves time, thereby allowing them to focus on income-earning activities. Participants were also found to engage in high-risk behaviours, including unsafe sex, which increased their risk of contracting HIV/AIDS and other sexually transmitted infections.

Introduction

1.1 Background: The Phenomenon of Street Children

Millions of children and adolescents live and/or work on the streets of large cities worldwide¹. Many of these children are denied access to their basic rights, including education, healthcare, and protection from abuse and exploitation. On the streets, they frequently survive by scavenging, begging, stealing or working in informal sectors in low-paying jobs. Some exchange sex for money, thus increasing their risk of contracting HIV and other sexually transmitted infections.²

In sub-Saharan Africa, the number of street children has been increasing due to the growing number of AIDS orphans. UNICEF (2003) estimated that eight out of every ten children who have lost parents to HIV/AIDS live in sub-Saharan Africa, and between 1990 and 2001 the proportion of orphans whose parents died from HIV/AIDS rose from 3.5% to 32%. UNAIDS (2006) estimated that there are more than 34 million orphans in the region, 11 million of them orphaned by HIV/AIDS. In the past, most orphaned children were cared for within the extended family.³ However, due to the high death rate associated with AIDS and increasing poverty in sub-Saharan Africa, many orphans live in elderly- or child-headed households that are unable to meet the children's basic needs of food, clothing and shelter.⁴ As a result, many orphans have no option but to live and work on the streets.

In the Tanzanian context, visible numbers of street children have been apparent since the 1990s.⁵ Various factors have led to this increase. As in other parts of sub-Saharan Africa, the AIDS epidemic is one of the major factors. According to UNAIDS, about 1.1 million Tanzanian children under 17 years of age have lost one or both parents to AIDS.⁶ As parents fall sick and die of AIDS, many families fail to take care of the orphans. As a result, many children must take on the responsibilities of parents and/or guardians, including income generation to support their households.

The impact of AIDS has been compounded by high poverty levels. As a result of growing economic disparities between urban and rural areas in Tanzania,⁷ many people from rural areas migrate to urban areas in search of economic opportunities. Due to economic hardship, many parents/guardians in urban areas also send their children to the streets to engage in income-generating activities, and eventually these youth start living on the streets. A number of studies have found that extreme poverty in many families lay at the centre of the increasing number of street children in Tanzania.⁸ For instance, Evans (2002) found that the inability of parents and guardians to pay school expenses resulted in high drop-outs rates which, in turn, led to children having to earn a livelihood on the street.

Non-income factors such as marriage breakdown and divorce, young parenting and domestic violence are also contributors to the increase in the number of street children in Tanzania. Research by the Mkombozi Centre found that family breakdowns and the inability

1 UNICEF 1989, Raffael et al. 1993, Scanlon et al. 1998

2 Raffael et al. 1993, Evans 2002, Anarfi 1997, Kamala et al. 2001

3 Ntozi 1997, Barnett & Whiteside 2002

4 UNICEF 2003, Loewenson & Whiteside 2001

5 Kopoka 2000, UNICEF 1990

6 UNAIDS 2006

7 Messkoub 1996, Cornia et al. 1987, Global Exchange 2001

8 Lugalla & Mbwambo 1999, Evans 2002, Community Development Department 2002, Yamamoto 1996

of extended families to take care of the children are major factors in the increase in children living on the streets in Kilimanjaro.⁹ Moreover, ILO/IPEC (2001) found that many children migrate to the streets in search of employment because of physical abuse and domestic violence, including sexual abuse and exploitation by relatives, parents and/or guardians.

In addition, secondary schools and vocational training centres in Tanzania do not have sufficient places for all children who complete primary education. For a large proportion of Tanzanian children, their formal education finishes at the end of primary school. As a result, many of these children end up on the streets. A review by the Social Action Trust Fund (SATF), a non-government organisation supporting orphans and vulnerable children (OVC) in Tanzania, shows that out of the 1,447 OVC surveyed who completed Standard VII in 2004-2005, only 311 children (21% of the total) were selected to join government secondary schools. Left without support, many of these children have no alternative but to earn a livelihood on the street.

The Ministry of Community Development, Gender and Children (MCDGC) has put in place a child development policy, which aims to educate the public about the rights of children, and to provide guidance on caring, supporting and protecting children in Tanzania.¹⁰ The policy, however, does not provide specific guidance on supporting vulnerable groups of children, such as children living on the streets, which necessarily impacts how interventions are designed for these groups.

Of the many vulnerable members of Tanzanian society, children living on the streets are among the most at risk.¹¹ Street life exposes children to a variety of socio-economic problems, including ill health and injury.¹² Health policies, programmes and services are therefore required to meet the needs of street children. To improve the design and implementation of these interventions, it is important to understand the current circumstances of street children in Tanzania, their vulnerability to poor health outcomes, and the strategies they use to cope with illness.

1.2 Research Problem and Significance of the Study

Children who are living on the streets are vulnerable to social and economic problems which may not be experienced by other young people. In particular, life on the street exposes children to illness and disease.¹³ Ensuring street children's access to health services is crucial to improve their health and well-being. For this aim to be realised it is important to investigate the coping strategies used by street children when they get sick. This will help to design healthcare interventions and service delivery for this population.

In Tanzania, studies have investigated the vulnerability of street children to HIV,¹⁴ and a situation analysis of street girls has been conducted.¹⁵ The extent of child labour in Tanzania

9 Mkombozi Centre 2005

10 MCDGC 1996

11 UNICEF 1989

12 Raffael et al. 1993, Population Reports 2001, Swart-Kruger & Richter 1997, Anarfi 1997, Kabondo et al. 2004, Kamala et al 2001, Kadonya et al. 2002

13 WHO 1993

14 Evans 2002, Amury 2004, Rajani & Kundrati 1994

15 Yamamoto 1996

has also been examined.¹⁶ In addition, a number of studies have investigated the nature and causes of street children and the problems they face on the streets.¹⁷ No studies, however, have been conducted to assess coping strategies used by street children when they get sick. This has hindered the design of relevant health services and health programmes for street children in Tanzania, particularly in Dar es Salaam, which has a large population of children living and working on the streets.

This study seeks to provide information for the proper planning of health services and programmes to meet the special needs of street children. In particular, it aims to understand gender differences in health-seeking behaviours, so as to inform the design of gender-specific health interventions for both street boys and girls. The circumstances of street children must be well understood to improve their access to health services which, in turn, will contribute to improved health outcomes for this vulnerable population.

1.3 Research Objectives

The broad objective of this study is to understand the coping strategies used by street children when they get sick. More specifically, the study aims to:

- Identify the diseases or infections that children living on the streets frequently succumb to
- Determine the health-seeking behaviours of street children
- Identify the barriers to accessing health services faced by children living on the streets
- Provide recommendations on health services and programmes to meet the health needs of this population.

1.4 Research Questions

Information and data collection was guided by the following main research questions:

1. What diseases or infections are common to street children?
2. What behaviours put street children at risk of contracting infections and diseases?
3. How do street children seek treatment when they are sick?
4. Is there any significant difference in health-seeking behaviours between girls and boys?
5. Do street children use available health services? If so, why?

¹⁶ ILO/IPEC 2001

¹⁷ Lugalla et al. 1999, Mvungi 2002, Lugalla & Kibassa 2003

Literature Review

This chapter analyses literature relevant to street children and their vulnerability to disease. To begin, Section 2.1 provides the definition of children living on the streets adopted by the current study. Section 2.2 discusses Tanzania's national health policy with a focus on access to healthcare. Section 2.3 then discusses recent research on health-seeking behaviours of street children and their access to health services. The final two sections of the chapter examine evidence on the vulnerability of children living on the streets to disease (Section 2.4) and the gender dimension of street children's vulnerability (Section 2.5)

2.1 Definition of Children Living on the Streets

This study defines 'children living on the streets' as children 18 years or younger who live independently on the streets in urban areas, largely supporting themselves by begging and scavenging on the streets or working in the informal sector, without adult supervision and with little or no family contact. Street children may also be breadwinners for poor households or children from street families, i.e. those who were born on the streets.

2.2 Tanzania's National Health Policy and its Implications for Access to Healthcare

The overall objective of Tanzania's National Health Policy is to improve the health and well-being of all Tanzanians, with a focus on those most at risk, and to encourage the health system to be more responsive to the needs of the people.¹⁸ The policy's specific objectives include ensuring the availability and accessibility of health services to all people in urban and rural areas, and to sensitise the community to common, preventable health problems. It also seeks to improve the capability – at all levels of society – to assess and analyse problems and to design appropriate actions through genuine community involvement.

To better respond to the health needs of Tanzanians, the Health Sector Reform Programme was commenced in 1994. The major aim of the reforms was to develop a health service system that would be equitable, sustainable, and involve both public and private modalities of healthcare delivery. Under the reforms, the Government is committed to provide essential health service functions commensurate with its budgetary capacity, and to facilitate the development of quality services in both public and private sectors.

Significant progress in the health sector was achieved through the reforms. Health infrastructure has been expanded such that 93% of the population lives within 10 km of a health facility and 72% within 5 km.¹⁹ However, national averages do not take into account geographical and other local variations, quality of services, and direct and indirect costs which determine accessibility to services.

This is particularly important in the case of street children, as the majority of them have limited access to mainstream social services. Rajani et al. (2001) found that, despite the fact that public health facilities were very widely distributed and geographically accessible, cultural barriers and inattentiveness to adolescents' needs rendered health services inaccessible to the adolescent population, including street children.

18 MOH 1990

19 MOH 1996

Indeed, none of the objectives in the National Health Policy address adolescent health generally or street children specifically. Moreover, the policy recognises the need to improve and develop urban health services, but specific provisions are not included to meet the challenge of the large influx of street children into urban centres. There is little evidence of any awareness of street children as a population group with special needs. Of serious concern, the Government's guidelines regarding user charges for health services do not identify street children as a vulnerable group.²⁰ According to the guidelines, only children under five years of age are exempt from payment of fees. Street children are not identified as a special group to be exempt.

2.3 Accessibility of Healthcare and Health-seeking Behaviour of Children Living on the Streets

Evidence indicates that children who are living on the streets tend to under-utilise the existing health services.²¹ A number of factors contribute to this problem. Of importance is that the majority of health services and programmes have been developed by adults for adults.²² They often fail recognize the unique issues of adolescents in general, and children living on the streets in particular, and rarely accommodate their behaviours.

For example, Tanzania's main health programmes – such as maternal and child health, family planning and AIDS control – do not have a component for street children. The principal focus of these programmes is the reproductive role of women.²³ The programmes also target children in school but commonly exclude youth who have dropped out and street children. Yet children living on the streets are amongst the poorest and most vulnerable adolescents.

It is also important to note that for children living on the streets the struggle to get daily food and shelter overrides concerns about their health. Kruger and Ritcher (2003) found that street children are more oriented to the present and not concerned with their future health. Other studies have found that even if these children appreciate the risk of getting infected with diseases, many believe that they are personally invulnerable.²⁴ This perception may result in children not using health services.

Even if street children want to use the health services, many are below the age of consent but neither have a parent/guardian to accompany them for treatment nor the necessary documentation.²⁵

The inaccessibility of healthcare to street children in sub-Saharan Africa has been exacerbated by the introduction of user fees for health services since the mid-1980s.²⁶

20 MOH 1997a

21 Kabondo et al 2004, Anarfi 1997, Kruger & Ritcher 2003, Dilip 2004

22 WHO 1993, Rajani et al. 2001

23 MOH 1997b, MOH 1994, TACAIDS 2003

24 Anarfi & Antwi 1995, Amury 2004

25 WHO 1993

26 Mujinja 2000

In South Africa, children living on the streets said that they did not seek treatment when they were ill because they did not have money and instead tried to sleep off their illnesses.²⁷ In Tanzania, an unpublished study showed that when street children fall ill they have to look after themselves, prescribe their own medications, and buy drugs from local pharmacies.²⁸ Research in Tanzania and Ghana also found evidence of repeated illness and sexually transmitted infections (STIs) among street children that were inadequately treated.²⁹ Left untreated, STIs can increase susceptibility to further infections, including HIV/AIDS.

2.4 Vulnerability of Street Children to Diseases and Illnesses

Children living on the streets worldwide are vulnerable to a range of health and other problems, including HIV/AIDS, which may not be experienced by other young people.³⁰ A number of behavioural and social characteristics of street children substantially increase their risk of contracting HIV. Studies have found that this population becomes sexually active earlier than most other groups of children, and they are known to engage in sexual activity with peers and adults from within and outside their social circle.³¹

Research has also shown that children living on the streets are likely to be raped or forced into sexual relationships on the streets, and that they use condoms infrequently and inconsistently.³² In addition, children living on the streets are more likely to exchange sex for money.

Furthermore, children living on the streets often receive limited information about reproductive health and HIV/AIDS due to illiteracy and non-attendance at school.³³ They also know very little about other STIs.³⁴ Akwara et al. (2003) argue that exposure to AIDS information tends to lead to higher awareness of the disease which, in turn, influences self-assessment of risk and a decrease in high-risk sexual behaviour, thereby reducing vulnerability to HIV.

In many societies, children living on the streets are also perceived as a threat. They are often vulnerable to discrimination and can be a very difficult population to reach and work with.³⁵ As a means of coping, many children living on the streets are involved in substance abuse. A study in Dar es Salaam found that street children were involved in marijuana smoking and substance abuse to endure tough jobs, discrimination and long days without food.³⁶ Drug use may further lead to a high risk of psychiatric disorders, such as mood disorders, depression, suicide, anxiety and phobias, sleep disorders and eating disorders.³⁷

The environments in which street children live can make them emotionally and physically vulnerable. In Tanzania, these children commonly sleep in hazardous environments such as in abandoned, broken-down vehicles, at bus stops and in bars.³⁸

27 Kruger et al. 2003

28 Amury 2004, Kamala et al. 2001

29 Kamala et al. 2001, Anarfi et al. 1995

30 Situmorang 2005, Kamala et al. 2001, Rajani et al. 2001

31 Raffael et al. 1993, Swart-Kruger et al. 1997, Rotheram-Borus et al. 1991, Anarfi 1999, Rajani and Kundrati 1994, Rau 2002

32 Kruger et al. 2003, Anarfi 1997, Anarfi 1999, Kamala et al. 2001, Rajani et al. 1994

33 Filgueiras 1993, Swart-Kruger et al. 1997, Campbell & Ntsabane 1995

34 Anarfi 1997

35 Csete & Whitman 2001, Ntarangwi 2003

36 Kamala et al. 2001

37 Friedman 1989, WHO 1993

38 Lugalla & Kibassa 2002, Mvungi 2002, Kopoka 2000

In addition, many children on the streets live, work and scavenge in dangerous and/or unhygienic conditions, exposed to discarded hospital needles and syringes, broken bottles and dangerous chemicals. They typically wear no protective gear such as gloves, helmets, gum boots, masks and glasses.³⁹ Living in such conditions at a young age subjects children to considerable stress and emotional turmoil, and many of the strategies they use to cope with their problems are harmful to their health and increase their vulnerability to diseases. A World Health Organisation (WHO) study found that street children suffered health problems including injuries from accidents and violence as well as industrial and environmental poisoning.⁴⁰ In Mwanza, Tanzania, many working children and children living on the streets were engaged in quarrying, garage work, fishing and fish processing, and complained of body pains, injuries and abuse, as well as frequent respiratory illnesses, and water-borne and other water-related diseases.⁴¹

In addition, the unhygienic conditions on the streets expose children to infectious diseases such as diarrhoea, typhoid, cholera, tuberculosis, skin diseases, respiratory tract infections and parasitic infestations.

2.5 The Gender Dimension of Vulnerability

Girls who are living on the streets are even more vulnerable than boys. In addition to the health risks experienced by street boys, adolescent girls are at risk of early, unplanned pregnancies with minimal, if any, antenatal care. Severe health risks, including maternal mortality, are associated with practices for terminating pregnancies.⁴²

Furthermore, more girls than boys experience sexual abuse and exploitation in the streets. In Zimbabwe, a study of sexually active street girls found that the majority had been forced into sex⁴³ and, in Ethiopia, a study involving 32 girls living on the streets reported that 21 of the participants had been raped.⁴⁴ A Tanzanian study by ILO/IPEC showed that most girls working on the streets had experienced sexual abuse.⁴⁵ Similarly, Evans (2002: 59) found that:

A street girl is in a lot more danger than a boy. Many, many women at the bus stand are raped. You hear the older boys saying, 'There are girls sleeping in a certain place, let's go and find them'. But a boy can sleep anywhere; he does not have any problem because he is a boy.

However, younger boys living on the streets are also particularly vulnerable to sexual exploitation. A focus group discussion with street children in Mbeya Municipality elicited that younger boys were engaged in sex for protection while others were raped by older men or older boys in the streets.⁴⁶ In South Africa, younger boys living on the streets claimed to have been often raped by older boys.⁴⁷ These factors significantly increase the risk of contracting HIV/AIDS and other STDs.

39 Kadonya et al. 2002, Lugalla et al. 2002, Mvungi 2002, Kopoka 2000

40 WHO 1993

41 Kadonya et al. 2002

42 WHO 1993, Kabondo et al. 2004, Kamala et al. 2001

43 UNICEF 2001

44 Lalor et al. 1992

45 ILO/IPEC 2001

46 Amury 2004

47 Kruger et al. 2003



Methodology

This chapter describes the methodology used for this study. Section 3.1 outlines the research approach, and Section 3.2 identifies the study locations. The sampling procedure is then discussed in Section 3.3 followed by description of data collection procedures and instruments in Section 3.4 and data analysis in Section 3.5. The final two sections of the chapter discuss the ethical considerations involved in conducting the research with children (Section 3.6) and the limitations of the study (Section 3.7).

3.1 Research Approach

Denzin (1989) argues that triangulation of research methods can overcome personal biases and limitations that stem from the use of a single research method. Therefore, both quantitative and qualitative data collection instruments were used in the current study. To begin, individual interviews using a structured questionnaire were conducted to gather quantitative data on the coping strategies used by children living on the streets in case of illness, as well as demographic data of participants. Qualitative methods were then conducted to verify the data collected and to gain an in-depth understanding of the children's coping strategies. The qualitative methods employed were semi-structured interviews, and participatory mapping and drawing exercises. To facilitate comprehensive responses to questions related to diseases or infections experienced by participants, a list of diseases common to street children was employed. However, no medical tests were conducted to diagnose the diseases identified by participants.

3.2 Study Locations

The study was conducted in the three municipal districts of Dar es Salaam: Ilala, Kinondoni and Temeke. The city was selected for the research because it is the largest urban area in Tanzania, and attracts many children and youth who are searching for employment.⁴⁸ In the most recent national census, Dar es Salaam's population was estimated to be 2,497,940 people.⁴⁹ The city has a high population of street children who are engaged in the informal sector.

Data was collected from the following sites:

- Ilala District – sites included Kariakoo market area, Mnazi Mmoja, Kidongo Chekundu, Buguruni, Feri, Fire and Jangwani;
- Kinondoni District – sites included Ubungo terminal bus stand, Mwenge, Magomeni and Manzese (Uwanja wa Fisi);
- Temeke District – sites included Chang'ombe, Tandika Sokoni, Kurasini (dampo area) and Mbagala.

The research took place in areas where street children typically congregate, such as bars, bus stands, local brew shops, shopping centres, market areas and on the streets.

⁴⁸ UNICEF 1990

⁴⁹ United Republic of Tanzania (URT) 2002

3.3 Study Participants and Sampling Procedure

As defined in Section 2.1, the study's target population was children who were living or working independently on the street. Owing to the lack of census data on children living on the streets in Dar es Salaam and the difficulty in tracking this highly mobile population, the study established the following criteria for selecting respondents:

- i) children had to be between 10 and 18 years of age; and
- ii) had to have lived/worked on the street for more than three months.

In total, 272 children participated. Survey interviews were conducted with 234 participants, participatory exercises engaged 30 participants (six groups of five children), and semi-structured interviews involved 8 participants (see Table 1).

Table 1: Number of Participants

Methods	Number of Participants
Survey interviews	234
Participatory exercise	30
Semi-structured interviews	8
Total	272

Control Sample

To ensure that study findings were reliable, additional interviews were conducted with a control sample of 40 boys who had previously worked or lived on the streets. The interviews involved 20 children from Dogodogo Children Centre in Kigogo (Ilala District) and Children in the Sun Centre in Mbezi (Kinondoni District). The children interviewed had been staying in the centres for more than one year. Participants were asked to describe their experiences while they were on the street. The same questionnaire used in the field was administered to the control sample. Findings from the control sample were compared to data collected in field interviews to assess any bias in the responses of children currently living and working on the streets (see Appendix 1).

3.4 Survey Instruments and Data Collection

Structured interview questionnaire

The structured interview questionnaire was developed in English and translated into Swahili. (see Appendix 2 for the questionnaire in Swahili and Appendix 3 for the English version). Principal topics covered by the questionnaire included:

- Social and demographic characteristics of participants, including age, gender, place of residence and level of education
- Diseases experienced by participants
- Coping strategies used by street children in case of illness
- Health-seeking attitudes and behaviour, including those related to HIV/AIDS.

It has been argued that information obtained using structured questionnaires is limited by the categories imposed by the researcher, and interesting replies by participants may be lost.⁵⁰ To minimise this problem, an 'other' category was provided for every question to ensure that all responses from the children were able to be recorded.

The questionnaire was piloted with 25 street children (who were not included in the main sample of 234 children interviewed). Based on pilot findings, revisions were made to the questionnaire. For example, it was found that children had difficulty recalling illnesses they had suffered over the past six months and, hence, this period of time was reduced to three months in the final questionnaire.

Data collection was face-to-face. Interviews were conducted in Swahili on the streets at research sites after respondents had been given information about the research and had provided their oral consent.

Semi-structured interviews

Semi-structured interviews were conducted to further investigate the coping strategies and health-seeking behaviour of street children. The researchers followed an interview guide (see Appendix 4), but also probed for more information based on participants' responses. The interviewer also had a chance to clarify responses from the interviewee. Moreover, the semi-structured interviews provided greater insight into the social factors underlying street children's coping strategies and health-seeking behaviour. The researchers piloted the interview guide with two children prior to field interviews, but no changes were needed to the guide.

Field interviews were conducted on the streets at research sites. Each interview lasted for about 30-40 minutes, and was tape-recorded with the permission of participants.

Participatory exercises: Mapping and drawing

Johnson and Nurrick (2000) note that:

Participatory visual methods can play an important role in helping children to present and analyse their own perspectives on issues that directly affect their lives. Such an approach is shown to promote learning and education (particularly in the non-formal sector) and also to provide the means by which children can inform decision-makers of their priorities and concerns, and provide solutions to problems that face them.

In the current study, mapping and drawing exercises were used to allow more active participation of children in sharing information about their coping strategies. The exercises did not require reading or too much writing but rather involved drawing, mapping and discussions. Large quantities of information were produced as discussions during the exercises were tape-recorded.

Participatory exercises were conducted in Swahili, on the pavements, and involved six groups of five street children each (see Appendix 5).

50 Bryman 2001, Memon 1990, Denscombe 2003, Franfort-Nachmians and Nachmians 1996

3.5 Data Analysis

After data collection, open-ended questions were pre-coded. Preliminary analysis was also completed on the demographic characteristics of participants. A second stage of analysis was then completed using the *Statistical Package for Social Science* (SPSS) and chi-square tests were performed to determine any significant differences between genders in relation to children's coping strategies and health-seeking behaviours.

Qualitative data was analysed in accordance with analytical procedures suggested by Marshall and Rossman (1999). Recordings from the semi-structured interviews were transcribed by listening to the tapes in conjunction with reading fieldwork notes. This exercise was done separately by two people (the researcher and a research assistant) and transcriptions were compared to ensure reliability of the findings. Transcriptions were then translated from Swahili to English, after which the common themes and patterns were identified.

3.6 Ethical Considerations

The ethical principle governing research is that participants should not be harmed as result of the research, and should give their informed consent.⁵¹ In particular, Allmark (2002) notes that research with children must be ethically sound, and undertaken with proper respect for the children's safety, integrity and privacy. It is also argued that consent to participate in the research should be given by someone competent to do so.⁵²

However, in the current study, parents/guardians were not available, so children were asked personally for their consent to participate in the research. All participants were given full information regarding the purpose of the research, what was expected from them, how long the interviews or exercises were expected to last, and that participation was voluntary. In addition, the research did not expose the children to any physical or emotional stress.

3.7 Limitations of the Study

There is potential for interviewer bias during face-to-face interviews. However, the use of a standardised questionnaire as well as piloting of the questionnaire, and training of interviewers on interview skills helped minimise this effect.

Some children were also scared to be interviewed by strangers (researchers) and at times were reticent to discuss their experiences, especially personal behaviours. Sound interviewing skills, such as establishing rapport, putting children at ease and remaining non-judgemental⁵³ were used to help children overcome their shyness. In addition, children were involved as facilitators in the mapping and drawing exercises, which encouraged their active participation.

51 Bowling 2002

52 Beuchamp and Childress 2001

53 Bowling 2002

Privacy was another problem as the study was conducted on the streets at research sites. There were some distractions from other street children and onlookers during interviews. Therefore, where possible, depending on the flexibility of individual respondents, interviews were conducted in relatively quiet places, away from the busy streets where most street children were found. Of note, girls were invisible during daytime and many refused to be interviewed in areas where there was no privacy. However, conducting interviews early in the morning and late in the evening in the places where they sleep (street pavements) helped to increase the sample size of street girls who participated.

Six participatory exercises were conducted instead of the planned eight. Participants for the exercises in Temeke District did not turn up and, due to the time limitation in the research clearance letter for that district, it was not possible to re-schedule this fieldwork.

In addition, some street children were unable to be interviewed as they were drunk at the time of interview. This was a frequent problem in Manzese (*Uwanja wa Fisi*), where most children in the area were drunk. Nevertheless, researchers persevered and interviewed those respondents who were ready and available.

Findings and Discussion

This chapter presents and discusses the findings of the study. Section 4.1 provides the general characteristics of respondents. Data on diseases and illnesses that street children frequently suffer, and their health-seeking behaviours are then presented in Sections 4.2 and 4.3 with a focus on gender differences in coping strategies. Section 4.4 details children's reasons for not using health services, and Section 4.5 presents children's suggestions to improve their access to healthcare. Finally, Sections 4.6 and 4.7 outline and discuss findings on behaviours and activities that put street children at risk of contracting diseases and illnesses, including HIV/AIDS.

4.1 General Characteristics of Respondents

Structured interviews were conducted with 234 street children, of whom 140 (59.8%) were boys and 94 (40.2%) were girls. Semi-structured interviews and participatory exercises involved 38 respondents (23 boys and 15 girls). Table 2 summarises data on the characteristics of respondent involved in structured interviews including age, length of time on the street, level of education attained and usual sleeping place.

Almost two-thirds of the street boys (63%) were aged between 11 and 15 years, and approximately one-quarter (24%) were aged between 16 and 18 years. Only 13% of respondents were aged 10 years. Among the street girls interviewed, 33% respondents were aged between 11 and 15 years, while 61% were aged between 16 and 18 years.

These findings indicate that boys leave home earlier than girls, which may be attributed to prevailing gender roles in many ethnic groups in Tanzania. Whilst girls are commonly socialised for reproductive roles⁵⁴ and obliged to engage in household chores, their male counterparts are frequently socialised to engage in remunerative productive roles, thus allowed to go on the streets to look for money earlier than girls.^{55,56} These findings are consistent with a study in Mbeya Municipality which found that boys are likely to start living on the streets at a younger age than girls.⁵⁷

Table 2: General Characteristics of Respondents

Variable	Male (N=140)		Female (N=94)	
	Frequency	Percentage	Frequency	Percentage
Age				
10 years	18	12.9	6	6.4
11-15 years	88	62.9	31	33.0
16-18 years	34	24.2	57	60.6
How long living / working in the street (years)				
Under one year	28	20.0	28	29.8
One year	26	18.6	15	16.0
More than one year	86	61.4	51	54.2

54 Reproductive roles involve the care and maintenance of the household and its members, including bearing and caring for children, preparing food, collecting water and fuel, housekeeping and family health care. It is almost always the responsibility of women and girls (Moser 1993).

55 Productive roles involve production of goods and services usually paid or income generating (Moser 1993).

56 Mwangeni et al. 2001

57 Amury, 2004

Table 2 (continued): General Characteristics of Respondents

Variable	Male (N=140)		Female (N=94)	
	Frequency	Percentage	Frequency	Percentage
Education Level				
No education	26	18.6	15	16.0
Drop-outs	70	50.0	30	31.9
In school	4	2.8	6	6.3
Completed primary education	40	28.6	43	45.7
Place to Sleep				
On the street	48	34.3	18	19.1
At home	41	29.3	51	54.3
Ghetto	51	36.4	25	26.6

The findings further show that over half of street boys and girls (61% and 54% respectively) had lived and/or worked on the streets for more than one year. This suggests that many of these children started living on the streets at very young ages. The children who were working were commonly engaged in activities such as selling foods, fruit or sweets, washing cars, or involved in illicit activities, such as drug dealing and commercial sex work.

With respect to educational attainment, half of the street boys (50%) and almost one-third (32%) of girls had dropped out of school. This is consistent with a study done by Mkombozi Centre (2005) in Kilimanjaro, which found that many children living on the streets were school drop-outs. Many parents or guardians are too poor to provide school requirements for their children, such as uniforms and books. As a result, many children drop out of schools and find their way to the streets.

In addition, more street girls (46%) than street boys (29%) had completed primary school. In this respect, the findings do not match those of Anarfi and Antwi (1995) whose study in Ghana found that street boys were more likely to have basic primary education than street girls. On other hand, 18% of street boys and 16% of street girls have never attended school. It may be argued that because street boys in the current study were living on the streets earlier than street girls, that they were less likely to finish or to attend primary school.

Furthermore, more street boys (34%) than street girls (19%) are sleeping on the streets while 36% of street boys and 27% of street girls are sleeping in a 'ghetto'. According to them, a ghetto is a cheaply rented room which accommodates between 10-20 children and is mainly rented for sleeping at night. This type of sleeping accommodation arrangement may expose children to the danger of getting air borne diseases.

4.2 Common Diseases and Illnesses of Street Children

In order to assess the health-seeking behaviour and coping strategies used when street children fall ill, it was important to know if respondents had ever been sick. Therefore, participants were asked whether they had been sick during the past three months and to describe the diseases they had suffered. The findings are summarised in Tables 3 and 4.

Table 3: Children Who Experienced Sickness During the Last Three Months

Response	Male (N=140)		Female (N=94)	
	Frequency	Percentage	Frequency	Percentage
Yes	132	94.3	84	89.4
No	8	5.7	10	10.6

Results in Table 3 show no significant difference ($p>0.05$) by gender in relation to whether or not children had been sick during the past three months. Almost all children had suffered from diseases or illnesses during the past three months (94% of street boys and 89% of street girls).

Table 4: Diseases/Illnesses Frequently Experienced by Street Children*

Type of Disease/Illness	Male (n=132)		Female (n=84)	
	Frequency	Percentage	Frequency	Percentage
Diarrhoea	33	25	24	28.6
Skin diseases ⁵⁸	53	40.2	26	30.9
Chest diseases ⁵⁹	33	25	14	16.7
Fever	87	65.9	58	69
Headache	39	29.5	36	42.9
Stomach ache	41	31.1	29	34.5
Accidents/Injury**	50	37.9	13	15.5

Notes: * This was a multiple response question, therefore, total frequencies and percentages exceed the total sample size (N=234) and 100% respectively, as some children suffered more than one disease during the last three months;

** $\chi^2 = 13.055 =$ significant difference ($p<0.01$) between genders

Data in Table 4 show that around two-thirds of street boys (66%) and street girls (69%) had suffered from fever. In addition, 40% of street boys and 31% of street girls had suffered from a skin disease during the past three months.

Results also show a significant difference ($p<0.01$) by gender in relation to experiencing accidents and injuries in the streets. Overall, 38% of boys had experienced injuries and accidents during the past three months, as compared to 16% of girls. This may be related to the nature of the work that boys are engaged in when on the streets. Kadonya et al. (2002) found that many street boys in Mwanza, Tanzania – who were involved in quarrying, garage work, fishing and fish-processing activities – complained of sustaining injuries.

Consistent with interview data, findings from the participatory group exercises revealed that street children are at risk of suffering diseases and illnesses. Figure 1 below illustrates the illnesses (*magonjwa*) suffered by participants when they were still living at home (*nyumbani*), and when they were living on the streets (*mtaani*).

58 Skin diseases referred to any type of skin rash

59 Chest diseases referred to pain in chest

No significant differences ($p>0.05$) by gender were found in relation to their health-seeking behaviour. Buying medicine at the local shop or pharmacy was the most common means of seeking treatment when they get sick (45% of boys and 46% of girls). This type of behaviour was also revealed in responses during the semi-structured interviews, as shown by the following excerpt:

'I don't usually go to the hospital when I am sick. Many times I bought drugs from pharmacies.' (Girl, 13 years)

Results were further confirmed by participants' input during the drawing exercises. Figure 2 illustrates a respondent going to the local pharmacy to buy medicine.

Figure 2:
A 14-year-old boy's
drawing of himself (left)
going to the pharmacy
(right) to buy medicine



Sharing medication was another means of coping with illnesses, as evidenced by one child:

'When I was suffering from malaria...I did not buy medicine. My friend gave me SP tablets⁶⁰ which were left over from when he was sick...After taking those tablets I got cured.' (Boy, 16 years)

By buying medicine direct from pharmacies or sharing with friends, street children are exposed to the danger of taking drugs without a proper diagnosis or prescription from a health practitioner. They may not receive the correct medication and recommended dosage, or complete the required course of treatment to fully recover. Inappropriate and/or incomplete treatment, may also contribute to the further development of drug resistance.

Data in Table 5 further reveal that 23% of boys and 27% of girls went to hospital when they were sick. However, almost one-quarter (24%) of boys and 18% of girls said that they sleep off their illnesses, which may seriously impact the children's health through delays in getting proper treatment. This behaviour is further evidenced by Figure 3, a drawing done by a 12-year-old boy, which shows him lying down when he was sick.

⁶⁰ Sulphadoxine-Pyrimethamine

Figure 3:
A street boy's drawing
(12 years old) of himself
when he was sick –
'I was suffering from
headache and slept
without medicine.'



Delay in seeking treatment and self-treatment has also been identified as a coping strategy used by the majority of poor people in Tanzania. Many tend to mobilise resources for seeking treatment only when they are seriously sick.⁶¹ This generalised behaviour may influence the coping strategies during illness used by children living on the streets.

It should also be noted that families with low incomes also tend to under-utilise available health services. Poor households are frequently forced to resort to self-treatment or ineffective alternatives, and may access facilities too late for care, often with fatal consequences. Many of these households resort to traditional healers and/or buying medicine from local shops or pharmacies.⁶² Children from low-income families may, therefore, be using the same health-seeking behaviours and coping strategies as children who are living on the street.

To better understand street children's behaviour, respondents were also asked to give their reasons for choosing the three coping strategies most commonly cited: 'going to a hospital', 'buying medicine' and 'sleeping off the illness'. The results are summarised in Tables 6 through 8.

Table 6 shows that out of the 50 respondents who chose to go to a hospital, 38% of children went because they were very sick, 30% because they trusted that the hospital services were effective, and 32% because they were advised to do so by their friends or relatives.

61 Save the Children 2003

62 Women's Dignity Project 2004, Guy 2003, Save the Children 2003

Table 6: Children’s Reasons for Going to the Hospital When Sick

Reason	Respondents (n=50)	
	Frequency	Percentage
Very sick	19	38
Hospital services are effective	15	30
Advised by a friend/relative	16	32

From these results, it may be argued that ‘going to hospital’ is actually not the first alternative for the 38% of children who reported going to hospital only when they became seriously ill. This is also supported by the following quote from a semi-structured interview:

‘If am not serious sick, I just sleep off my illness or I buy drugs from pharmacies. I go to the hospital when I am seriously sick.’ (Boy, 14 years)

Table 7: Children’s Reasons for Buying Medicine When Sick

Reason	Respondents (n=92)	
	Frequency	Percentage
Cheap	42	45.6
Saves time and nearby	26	28.3
Effective	24	26.1

The results in Table 7 show that 46% of children opted to buy medicine when they were sick because it was the cheap alternative, while 28% of respondents chose to buy medicine because this saved time and it was available nearby. In addition, findings from the semi-structured interviews revealed that street children spent most of their time looking for money to buy food, and hence they are not likely to spend much time seeking treatment. As one boy pointed out:

‘When I get sick I buy drugs from local pharmacies and I go to the war (begging)... You know, brother, if you don’t go vitani (in the war)...there is no food...That’s why I like going to the pharmacy because it saves my time.’ (Boy, 13 years)

Street children’s struggle to get food each day can, therefore, override any concern for their health. Amury (2004) also found that many street children were not concerned about their health because they had more pressing needs than seeking treatment.

Table 8: Children’s Reasons for Sleeping Off Their Illnesses

Reason	Respondents (n=46)	
	Frequency	Percentage
No alternative	27	58.7
Effective	7	15.2
Very sick	12	26.1

Data in Table 8 show that more than half of the respondents (59%) who chose to sleep off their illnesses said that they had no other alternatives, and 26% of respondents revealed that they sleep off their illnesses because they were so sick that they could not go to hospital and they did not have anyone to help them to the hospital.

To further assess their coping strategies, participants in the semi-structured interviews were asked about sources of cash for treatment. Several responses are presented below:

'When I suffered from gonorrhoea, I first used traditional medicine from Masai...but I did not get well...I [then] decided to buy medicine from pharmacy...My condition became worse...Then my friends contributed money and took me to the hospital.' (Boy, 17 years)

'When I get sick...I tell my friends...If they have money they buy drugs for me...If we don't have money ...I just sleep...You know we are very poor...and our life here in the streets is very hard.' (Boy, 16 years)

'One day my friend was very sick...He was unconscious...We contributed some money and took him to the hospital. Therefore, if we don't have money we don't go to the hospital...until we contribute or borrow money from each other.' (Boy, 18 years)

'In the past two months I got problem in throat...It was so big that I failed to eat...I was coughing a lot and I became very thin...My friends told me to go to Mdoe Street...There is a traditional practitioner who treat me for 3000/-. My friends contributed money for me.' (Boy, 18 years)

These excerpts reveal that street children have very limited individual resources to access health services, and rely on their friends to contribute money for drugs or treatment and to assist in getting to a hospital if they suffer a serious medical problem.

4.4 Children's Reasons for Not Using Health Services

In order to gain more insight on how to improve access and provision of healthcare services for street children, the respondents who did not go to a hospital were asked to give their reasons for not doing so. The data in Table 9 show that more than half of the street boys (60%) and street girls (53%) did not go to hospitals because it was expensive. These findings indicate that user fees for hospital services – that were introduced during the 1980s in Tanzania – are a frequent barrier to street children accessing health services. As one boy described:

'The main problem of getting treatment is money...Money to pay for medicine and money to pay to doctors...Money is the issue here.' (Boy, 15 years)

Table 9: Children’s Reasons for Not Going to the Hospital

Reason	Male (n=78)		Female (n=38)	
	Frequency	Percentage	Frequency	Percentage
Expensive	47	60.3	20	52.6
Hospital staff are unfriendly	10	12.8	11	28.9
To save time	11	14.1	2	5.3
Illness not serious	10	12.8	5	13.2

In addition, 13% of boys and 29% of girls did not go to hospital because they considered that hospital staff are unfriendly. As one child related:

‘Other watchmen in the hospital prevent us from entering the hospital gates. They say we are going to beg in the hospital.’ (Girl, 13 years)

Likewise, children’s illustrations from the group drawing exercises show that children living on the streets do not like to go to hospital because they are chased away by the hospital staff. Figure 4 shows a street girl being chased away by the guard, before she enters the hospital gate. The words – ‘You won’t get medicine here unless you have money’ – are written between her and the hospital.

Figure 4:
A drawing by a 15-year-old street girl (left) being chased away from the hospital by a watchman (right).



WHO (1993) also argued that street children are not likely to go to hospital because hospital staff treat them very badly.

4.5 Children’s Suggestions to Improve Healthcare Access

A principal aim of the research is to identify strategies to improve health outcomes for street children. Therefore, respondents were asked for their recommendations that would improve their access to health services. Findings are summarised in Table 10.

Table 10: Children's Suggestions to Improve Healthcare Access

Suggestion	Respondents (n=188)	
	Frequency	Percentage
Free services	92	48.9
Cost should be reduced	24	12.7
There should not be a long queue (i.e. reduce waiting time at the hospital)	30	15.9
Staff should be friendly	20	10.5
Don't know	32	17

Consistent with survey findings that the cost of treatment represented a significant barrier to participants in accessing care, almost half of the children (49%) suggested that health services should be free, and a further 13% felt that the cost should be reduced. In addition, 16% of respondents cited the need to reduce waiting time at the hospital. These recommendations again highlight children's requirements to save time and money so as to attend to their other immediate needs, such as food and shelter. The following quotes from the semi-structured interviews reinforce the importance of these issues:

'I would like the health services to be free of charge because even if you have 500/- it is not enough to get treatment... You need to buy a registration card [and] drugs... You need to pay 100/- for an injection... At the same time you don't have money to even buy food.' (Girl, 18 years)

'Health services should be offered free of charge to street children... Nurses and doctors should be compassionate to street children... When we go to the hospital ... they shouldn't be asking us too many questions... Even the police should give us a report that allows us to get treatment for free.' (Boy, 17 years)

4.6 Sexual Behaviours that Put Street Children at Risk of Contracting HIV/AIDS and other Sexually Transmitted Diseases (STDs)

To assess street children's risk of contracting HIV/AIDS and STDs, respondents were asked if they had ever had sex and, if so, whether they had ever used condoms. Findings are presented in Table 11.

Table 11: Children's Sexual Behaviour

Behaviour	Boys (n=139)		Girls (n=92)	
	Frequency	Percentage	Frequency	Percentage
Ever had sex				
Yes*	68	48.9	67	72.8
No	71	51.1	25	27.2
Ever used condom				
Yes	18/68	26.5	26/67	38.8
No	50/68	73.5	41/67	61.2

* $\chi^2 = 13.026 =$ significant difference ($p < 0.01$) between genders

Note: only 40 participants responded to sexual behaviour questions i.e. 24 boys and 16 girls.

A significant difference ($p < 0.01$) in sexual activity was found between boys and girls. Approximately three-quarters of the female respondents (73%) had ever had sex compared with about half of the male respondents (49%). However, it must be noted that the age of the girls surveyed was higher than the age of the boys. Overall, 61% of girls surveyed were aged 16 to 18 years compared with 24% of boys in this age bracket.

Among respondents who had ever had sex, 74% of boys and 61% of girls had never used a condom. Other studies have also found low levels of condom use among street children.⁶³ Evidence from the semi-structured also indicated that sexually active respondents used condoms infrequently or inconsistently, or were unable to negotiate safe sex. As one boy described:

'I prevent myself from getting HIV/AIDS by using condoms but I was infected with gonorrhea by my lover because I got drunk and I didn't use condoms...But I usually use a condom.' (Boys, 17 years)

One girl related feeling powerless to demand that her partners use condoms because they gave her money for living expenses:

'I would really like to protect myself against AIDS by using condoms...but when I tell my sexual partners to use condom they refuse...and I can't stop having sex with them because they give me money to buy food and clothes.' (Girl, 18 years)

This inability to negotiate safe sex among girls living on the streets significantly increases their risk of contracting HIV and other STIs.

Other research studies have found that women in general, not only street girls, lack confidence and communication skills in issues related to sexuality.^{64,65} Freudenthal (2001) also argued that young girls and women in many African countries not only lack the skills but also the resources and power to negotiate safer sex with men. This is particularly true for street girls who engage in sexual relations with older men or fellow street boys to get money to survive. Anarfi and Antwi (1995) found that girls on the street are more likely to be sexually exploited, but do not have the qualifications to secure an alternative livelihood.

4.7 Activities that Put Street Children at Risk of Contracting Diseases and Illnesses

Table 12 presents data on income-earning activities of respondents. Petty trade was the most frequently cited income-earning activity by both genders. About half of the boys (52%) and girls (48%) were engaged in petty trade. A significant difference by gender was found in the incidence of begging among street children. More girls (34%) were engaged in begging to earn income than boys (6%). Begging may expose street girls to sexual exploitation, hence increasing their risk of contracting HIV/AIDS. In contrast, a higher

⁶³ Campbell Ntsabane 1995, Anarfi and Antwi 1995

⁶⁴ Philemon & Kessy 2008

⁶⁵ Mwageni et al. 1998

proportion of boys (24%) than girls (18%) were engaged in scavenging. Scavenging is an extremely hazardous form of work, exposing children to infectious diseases, particularly hepatitis and tetanus. Small percentages of boys were also engaged in car washing and carrying luggage – 6% and 13% respectively – but no girls were involved in these two activities.

Survey findings are consistent with Kadonya et al. (2002) which found that street children were more likely to engage in petty trade, scavenging and begging.

Table 12: Income-earning Activities of Street Children

Activity	Male (n=140)		Female (n=94)	
	Frequency	Percentage	Frequency	Percentage
Petty trade	73	52.1	45	47.9
Scavenging	33	23.6	17	18.1
Begging*	8	5.7	32	34.0
Car washing	8	5.7	0	0
Carrying luggage	18	12.9	0	0

* $\chi^2 = 25.81 =$ significant difference ($p < 0.05$) between genders

With respect to income, street children revealed that they usually earn an average of Tshs 200/- to 1,000/- shillings per day, but money earned can fluctuate widely. As two children related:

'My life depends on begging...I usually get an average of 400/- a day. If it is a lucky day...I can even get 800/-.' (Boy, 14 years)

'Sometimes I get Tshs 1,000/- after scavenging...They usually buy used plastic bottles for Tshs 400/- per kilogram...so if I manage to get two kilograms I can get 1,000/- per day.' (Boy, 15 years).

As one girl pointed out children's income was not guaranteed especially if sick; sometimes they do not get any money at all.

'Sometimes it is very difficult to get even 100/-...especially when you are sick...You cannot work and hence you won't have money for food...We have a very difficult life.' (Girl, 16 years)

It is important to note that the minimum cost for an outpatient in government hospitals and dispensaries in Tanzania is Tshs 1,000.⁶⁶ If the study data reflect the level of income earned by street children – i.e., between Tshs 200-1,000 – then this clearly indicates that the cost of health service is unaffordable to children who are living on the streets.

The next chapter summarises the major findings and conclusions from the study, and provides recommendations on policies and interventions to reduce their risk of contracting diseases and to increase access to health services for children living on the streets.

⁶⁶ MOH 1997a

Summary of Key Findings, Conclusions and Recommendations

5.1 Summary of Key Findings

The major findings from this study are:

- Street children frequently experience illnesses, particularly fever, skin diseases, injuries, headaches, diarrhoea and stomach upsets.
- Children who are living on the streets under-utilise available health services.
- Buying medicine from local pharmacies and sleeping off illnesses are the major coping strategies used by street children when they are sick.
- No significant differences in health-seeking behaviour were found between boys and girls living on the streets.
- The cost of healthcare services, including user fees at health facilities, is the principal barrier deterring street children from accessing appropriate healthcare.
- Street children engage in high-risk behaviours, including unsafe sex, which increases their risk of contracting HIV/AIDS and STDs.
- The income-earning activities frequently cited by participants, especially scavenging and begging, exposed them to serious health and safety risks.

5.2 Conclusions

In light of study findings, a number of conclusions can be drawn. Firstly, the cost of health services for street children will need to be reduced if their access to healthcare is to be improved. The study showed that participants typically did not have the individual resources to pay the user fees and charges at health facilities, and so were unable to promptly access care when sick. However, providing free or reduced cost services *per se* may not be sufficient to achieve effective access. Healthcare staff must also be given training to better understand the special needs of street children and to provide youth-friendly services. In addition, the study revealed that under-utilisation of health services for street children is related to other factors such as time saving, so that they can attend to other immediate needs, such as food and shelter. Reducing waiting times at facilities will be needed to decrease the opportunity costs incurred by all patients, including street children. Achieving this outcome will require much greater investment in healthcare personnel.

Secondly, if street youth continue to engage in unsafe sex, they are at high risk of contracting HIV/AIDS. It is therefore critically important to provide children with information about prevention of HIV/AIDS. However, increasing their HIV knowledge and prevention skills *per se* may not mitigate their vulnerability to poor reproductive health outcomes. As the study revealed, their risk of becoming infected with HIV is closely related to the economic and social conditions which drive them to live and work on the streets. Therefore, efforts must be also made to improve their socio-economic opportunities and life skills – for example, through enrolment in school, vocational or entrepreneurial training, and support for livelihoods – if there is to be a significant and sustained impact on their vulnerability to HIV/AIDS. Organisations working with street children must also recognise and develop gender-sensitive interventions which address the multi-faceted social, economic and cultural factors affecting the vulnerability of street girls and street boys to HIV.

Lastly, a close relationship exists between abject poverty and street children. Improving the socio-economic status of poor households with children will be critical in reducing the numbers of children who have no alternative but to live and work on the streets as a means of survival.

5.3 Policy Recommendations

Tanzania's National Health Policy

The Tanzania National Health Policy should recognise the need to improve and develop urban health services to meet the challenge of the large influx of street children into urban centres and cities. Furthermore, the policy should recognise that street children are a population group with special needs. In addition, government guidelines regarding user charges for health services should take into account that street children are a vulnerable group that needs to be exempt from user fees. This is because the study reveals that children living on the streets are vulnerable to a range of diseases and other health problems, but under-utilise health services largely because the direct and opportunity costs of accessing health facilities are too high.

Tanzania's Child Development Policy

The Child Development Policy should identify street children as a special target group, and participatory strategies should be devised to reduce their vulnerability to diseases. Currently, the policy does not recognise street children as a target populations requiring special attention.

Tanzania's National AIDS Policy

The Tanzania Health Policy, the Tanzania National AIDS Policy, and the National Multi-Sectoral Strategic Framework on HIV/AIDS should take into consideration the involvement of street girls in commercial sex work and develop age-appropriate interventions. This study revealed that girls under 18 years of age are frequently involved in commercial sex work to make sufficient income to survive.

5.4 Recommendations for Practical Action to Support Street Children

Interventions to improve street children's access to healthcare should not only provide free health services to these children, but also address the social, cultural and environmental factors that restrict access. In this regard, the following interventions are recommended to Government, NGOs, and community- and faith-based organisations and individuals who are working with street children.

Basic Education and Vocational Training for Street Children

Efforts should be made to ensure that children who drop out of school and other street children are provided with basic primary education. Complementary Basic Education in Tanzania (COBET) may represent a key programme for this population.⁶⁷ COBET aims to increase access to education to out-of-school children, youth and illiterate adults, and ensure they have access to high quality learning opportunities. Special or non-formal primary education programmes will be critical in reaching children living on the streets.

⁶⁷ Ministry of Education and Culture (MOEC) 2002

It is acknowledged that the provision of street-side schools does not guarantee that street children will attend, as they may have more immediate needs such as securing food and shelter. In addition, street children may have developed health, psychosocial and behavioural problems in the streets that may affect their concentration and social relations at school.⁶⁸ Encouragingly, an intervention run by the the Faraja Trust Fund in Morogoro, Tanzania helped street children with their day-to-day survival concerns and, as a result, more children attended school consistently.⁶⁹ In addition, the Mkombozi Centre adopted the COBET programme in providing basic education to street children in Kilimanjaro region.⁷⁰ Using this approach, the Undugu Society of Kenya trained 2,000 street children between 1990 and 1995 who later joined ordinary primary and vocational schools.⁷¹ Based on this evidence, such a programme should be adopted and expanded to other regions and districts in Tanzania.

Ideally, children who have completed primary school should be offered tailor-made vocational training such as carpentry, masonry, cooking and tailoring as a foundation for income-generating activities. If possible, government and NGOs should also provide start-up tools to enable youth to establish small businesses. However, for street children to participate, vocational training will need to be cheap or free, conveniently located, and focused on practical skill⁷²; and should incorporate play, social or leisure activities and adopt a flexible schedule so as to motivate children.⁷³ With such training and support, street children will have much greater capacity to meet their basic needs.

Residential Homes and Drop-in Centres for Street Children

Residential homes and drop-in-centres are also required to provide shelter, food and health services to street children, as well as counselling. Services should be available to re-unite children with their families and communities where possible. However, children living on the streets cannot be forced to leave the streets. In order for the programme to be successful, this socialisation will likely take place in the streets, and the process of leaving the streets should be voluntary. The Faraja Trust Fund has been successful in re-uniting street children with their families using this approach.⁷⁴ But some children have completely lost contact with their families while others tend to return to the streets due to the hardship of life at home. In these cases, residential homes will be needed to provide long-term accommodation, care and vocational training until children reach adulthood. In addition, homes and drop-in centres should work in partnership with the families and communities from which the street children came. In Kenya, Kwetu Training Centre observed that the rate of street children who were returning to the streets after being re-united with their families fell by 50% after the programme started to involve parents and communities. These are important groups that need to be prepared for the integration of children returning to their homes.⁷⁵

Income-generating Projects for Children Living on the Streets

Older street children (16-18 years) should be helped to secure premises and facilitated to acquire start-up tools and capital to undertake income-generating activities. Training programmes in entrepreneurial skills would also be valuable. Car washing or shoe shine

⁶⁸ Volpi 2003

⁶⁹ Faraja Trust Fund 2000

⁷⁰ Mkombozi Centre 2005

⁷¹ Volpi 2003, Muli 2000

⁷² Frew 2003

⁷³ WHO 2000

⁷⁴ Faraja Trust Fund 2000

⁷⁵ Ruto 1999

shops are two examples of businesses that require little capital to start and can be conducted in the streets where children are living.

Partnerships Between Government, Health Facilities, Civil Society Organisations and Communities Involved in Programmes for Street Children

Increased collaboration between all stakeholders will help to maximise the benefits of interventions, minimise duplication of efforts, and utilise scarce resources more effectively.

5.5 Recommendations for Future Research

Greater knowledge and understanding of the circumstances of children living on the streets is necessary to inform appropriate policies, programmes and health services for this highly vulnerable population. The current study covered only one city, Dar es Salaam. A larger study encompassing different urban areas in Tanzania needs to be carried out. Such research would provide a more comprehensive understanding of the health-seeking behaviour and coping strategies used by street children in diverse urban contexts in Tanzania. A complementary assessment of existing policies and interventions to support street children will further facilitate the development of appropriate local and national strategies to meet the needs of all street children.

The study findings also indicate an urgent need to investigate street children's attitudes towards condom use and their ability to access them. Further research is also needed to examine the attitudes and practices of health personnel towards children living on the street. This will inform the development of a sensitisation and training programme for health workers to better understand the needs, behaviour and norms of this population, so that they are able to deliver appropriate youth-friendly services.

Bibliography and References

- Akwara A P, Madise J N and Hinde A. (2003). Perception of risk of HIV/AIDS and sexual behaviour in Kenya. *Journal of Biosocial Science*, 35: 385-411.
- Allison B, O'Sullivan T, Owen A, Rice J, Rothwell A and Saunders C. (1996). *Research skills for students*. London: De Montfort University.
- Allmark P. (2002). Ethical issues in research: The ethics of research with children *Nurse Researcher*, 10 (2). Accessed 20 June 2005 at http://www.nursing-standard.co.uk/nurseresearcher-archive/nr_archive/nrv10n2-2002/nrv10n2p0719.pdf
- Amury, Z. (2004). Street children's knowledge, attitudes and behaviours towards HIV/AIDS. Unpublished Masters Thesis, Queen Margaret University College, Edinburgh.
- Anarfi J K. (1997). Vulnerability to sexually transmitted diseases: Street children in Accra. *Health Transition Review*, 7 (Supplement): 281-306. Accessed 22 May 2005 at <http://htc.anu.edu.au/pdfs/Anarfi6.pdf>
- Anarfi J K. (1999). Initiating behavioural change among street-involved youth: findings from a youth clinic in Accra. In: Caldwell J C, Caldwell P and Anarfi J K (Eds) *Resistances to behavioural change to reduce HIV/AIDS infection in predominantly heterosexual epidemics in Third World countries*. Accessed 14 June 2005 at http://htc.anu.edu.au/pdfs/resistances_ch7.pdf
- Anarfi J K and Antwi P. (1995). Street youth in Accra city: Sexual networking in a high-risk environment and its implications for the spread of HIV/AIDS. *Health Transition Review*, 5 (Supplement): 131-151.
- Andersen R M. (1995). Revisiting the behavioral model and access to medical care: Does it matter? *Journal of Health and Social Behavior*, 36 (1): 1-10.
- Andersen R M and Newman J F. (1973). Societal and individual determinants of medical care utilization in United States. *Milbank Memorial Fund Quarterly*, 51: 95-124.
- Barnett T and Whiteside A. (2002). *AIDS in the Twenty-First Century: Disease and globalisation*. New York: Palgrave Macmillan.
- Beauchamp T and Childress J. (2001). *Principles of biomedical ethics*. 5th edition. Oxford: Oxford University Press.
- Bowling A. (2002). *Research methods in health: Investigating health and health services*. 2nd edition. Milton Keynes, UK: Open University Press.
- Bryman A. (2001). *Social research methods*. New York: Oxford University Press.
- Campbell K E and Ntsabane T. (1995). *Street children in Gaborone, Botswana: Causes and policy implications. Summary report No. 13*. Accessed 10 June 2005 at <http://www.uaps.org/Rapports/street.htm>
- Caron F, Gordin G, Otis J and Lambert L D. (2004). Evaluation of a theoretically based AIDS/STD peer education program on postponing sexual intercourse and on condom use among adolescents attending high school. *Health Education Research*, 19 (2): 185-197.
- Clay S and Bond V. (2004). *We didn't apply to be AIDS orphans*. Accessed on 5 June 2005 at <http://www.thezambian.com/Reports/aids.aspx>
- Community Development Department. (2002). Annual report, Mbeya Municipal Council Tanzania. Unpublished.
- Connell B. (2002). *Hegemonic masculinity*. In: Jackson S and Scott S. (Eds). *Gender: A sociological reader*. London: Routledge.
- Cornia G A, Jolly R and Stewart F. (Eds) (1987). *Adjustment with a human face: Protecting the vulnerable and promoting growth*. A study by UNICEF. New York: Oxford University Press.

- Csete J and Whitman L. (2001). In the shadow of death: HIV/AIDS and children's rights in Kenya. **Human Rights Watch**, 3 (4A). Accessed 20 May 2005 at <http://www.hrw.org/reports/2001/kenya/kenya0701.PDF>
- Denscombe M. (2003). **The good research guide for small-scale social research projects**. Milton Keynes, UK: Open University Press.
- Denzin K N. (1989). **The research act: A theoretical introduction to sociological methods**. 3rd edition. Englewood Cliffs, New Jersey: Prentice-Hall.
- Dilip A. (2004). Health-seeking behaviour among street children in Tanzania: A case study of Kinondoni and Ilala Districts, Dar es Salaam. Unpublished Masters Thesis, University of Dar es Salaam, Tanzania.
- Evans R. (2002). Poverty, HIV, and barriers to education: Street children's experiences in Tanzania. **Gender and Development**, 10 (3): 51-62.
- Ewald J, Shao I, Mhamba R, Mellander L, Narman A. (2004). **A matter of choice? Cost sharing in health and education from a rights of the child perspective**. Report prepared for a stakeholders meeting, 17 January 2004, Dar es Salaam: Goteborg University Centre for Africa Studies.
- Faraja Trust Fund. (2000). Street-side school report, Morogoro, Tanzania. Unpublished report.
- Filgueiras A. (1993). Defending children's rights: An AIDS prevention strategy. **AIDS Captions**, 1 (1): 10-13.
- Fisher A A, Laing E J, Stoeckel J E and Townsend J W. (1991). **Handbook for family planning operation research design**. 2nd edition. New York Population Council
- Franfort-Nachmians C and Nachmians A. (1996). **Research methods in the social sciences**. 5th edition. London: Arnold.
- Freudenthal S. (2001). **A review of social science research on HIV/AIDS SIDA/SAREC**. Accessed 22 May 2005 at http://www.somanet.org/youthproject_files/SAREC%20AIDS%20report.htm
- Friedman H. (1989). The health of adolescents: Beliefs and behaviour. **Social Science and Medicine**, 2 (3): 309-315.
- Fuglesang M. (1997). Lessons for life: Past and present modes of sexuality education in Tanzanian society. **Social Science and Medicine**, 44 (8): 1245-1254.
- Girbich C. (1999). **Qualitative research in health: An introduction**. London: SAGE Publications.
- Global Exchange. (2001). **How the International Monetary Fund and the World Bank undermine democracy and erode human rights: Five case studies**. Accessed 11 June 2005 at <http://www.globalexchange.org/campaigns/wbimf/imfwbReport2001.html>
- Guy H. (2003). **Improving the financial access of poor and vulnerable groups to health services. What strategy for Tanzania?** A briefing paper established under the SDC-STI SWAP Mandate 2003. Basel: Swiss Centre for International Health and Swiss Tropical Institute.
- Howard M M J. (1990). Helping teenagers to postpone sexual involvement. **Family Planning Perspectives**, 22 (1): 21-26.
- International Labour Organisation/International Programme on Elimination of Child Labour (ILO/IPEC). (2001). **Tanzania country report: 2000/2001 Integrated Labour Force and Child Labour Survey**. Accessed in May 2005 at <http://www.ilo.org/public/english/standards/ipec/simpoc/tanzania/report/rep2001.pdf>
- Johnson V and Nurrick R. (2000). Participatory visual aids with children. **Education Action 12**. Accessed 20 May 2005 at <http://217.206.205.24/resources/publications/EA12Eng/children.htm>

- Kabondo N, Nzima M, Msava L, Magalla A and Richardson J. (2004). **Orphans and vulnerable children in big cities: Considering special challenges for an intervention program in Dar es Salaam, Tanzania.** A paper presented at the XV International AIDS Conference, Bangkok, 2004. Accessed on 2 August 2005 at http://www.axios-group.com/en/library/posters/A0_13kabondo_26.pdf
- Kadonya C, Madihi M and Mtwana S. (2002). **Tanzania child labour in the informal sector: A rapid assessment.** A study for ILO/IPEC. Accessed 6 August 2005 at <http://www.ilo.org/public/english/standards/ipec/simpoc/tanzania/ra/infosec.pdf>
- Kaime-Atterhog W. (1996). Street children and prostitution: The situation in Kenya **News on Health Care in Developing Countries**, 10 (2): 27-32.
- Kamala E, Lusinde E, Millinga J and Mwaitula J. (2001). **Tanzania children in prostitution: A rapid assessment.** A study for the International Labour Organization International (ILO) Programme on the Elimination of Child Labour (IPEC). Accessed 4 August 2005 at <http://www.ilo.org/public/english/standards/ipec/simpoc/tanzania/ra/prost.pdf>
- Kindeberg T and Christensson B. (1994). Changing Swedish students' attitudes in relation to AIDS epidemic. **Health Education Research**, 9: 171-181.
- Kopoka A P. (2000). **The problem of street children in Africa: An ignored tragedy.** A paper presented at International Conference on Street Children and Street Children's Health in East Africa, Dar es Salaam, 19-21 April 2000. Accessed 12 April 2005 at <http://bij.hosting.kun.nl/esap/DAR/DAR11/General/Streetchildren.pdf>
- Kruger J M and Richter M L. (2003). South African street children at risk for AIDS? **Children, Youth and Environment**, 13 (1). Accessed 17 June 2005 at http://cye.colorado.edu/Vol13ArticleReprints/SAfricanStChldr_n_AfricanInsight.pdf
- Loewenson and Whiteside. (2001). **HIV/AIDS: Implications for poverty reduction.** United Nations Development Programme Policy Paper.
- Lalor K, Taylor M, Veale A, Ali H A and Bushra E. (1992). **Victimisation amongst children in Sudan and Ethiopia: A preliminary analysis.** Accessed 14 April 2004 at http://www.unicri.it/icvs/publications/pdf_files/understanding_files/26_VICTIMISATION%20AMONGST%20STREET%20CHILDREN.pdf
- Lugalla J P and Kibassa C G. (2002). **Poverty, AIDS and street children in East Africa.** Lewiston, New York: Mellen Press.
- Lugalla J P and Kibassa C G. (2003). **Urban life and street children's health: Children's accounts of urban hardship and violence in Tanzania.** Munster, Germany: Lit Verlag.
- Lugalla J P and Mwambo J K. (1999). Street children and street life in urban Tanzania: The culture of surviving and its implications in children's health. **International Journal of Urban and Regional Research**, 23 (2): 229-334.
- Memon A. (1990). Young people's knowledge, beliefs and attitudes about HIV/AIDS: A review of research. **Health Education Research**, 5: 327-335.
- Mellanby A R, Phelps F A, Crichton N J and Tripp J H. (1995). School sex education: An experimental programme with education and medical benefits. **British Medical Journal**, 311: 414-417.
- Messkoub M. (1992). **Deprivation and structural adjustments.** In: Wultys M (ed.) Development Policy and Public Action. Oxford: Oxford University Press.
- Messkoub M. (1996). The social impact of adjustment in Tanzania in the 1980s – Economic crisis and household survival strategies. **Internet Journal of African Studies**, 1. Accessed 20 April 2004 at <http://www.bradford.ac.uk/research/ijas/tazadj2.htm>

Ministry of Community Development, Women Affairs and Children (MCDWC). (1996). **Child Development Policy**. Dar es Salaam: United Republic of Tanzania. Ministry of Education and Culture. (2002). Non-formal education status report (Tanzania Mainland). Dar es Salaam: United Republic of Tanzania.

Ministry of Health (MOH). (1990). **Tanzania National Health Policy**. Accessed 20 June 2005 at <http://www.tzonline.org/pdf/Nationahealthpolicy.pdf>

Ministry of Health. (1994). **National Policy Guidelines for Family Planning Services, Delivery and Training**. Dar es Salaam: Government Printer.

Ministry of Health. (1996). **A Ministry of Health - Donor appraisal for Tanzania Mainland**. Dar es Salaam: United Republic of Tanzania.

Ministry of Health. (1997a). **Kamati ya Utekelezaji Utaratibu wa Wananchi Kuchangia Huduma za Afya: Mwongozo wa Utekelezaji wa Sera ya Wananchi Kuchangia Gharama za Afya katika Hospitali**. Dar es Salaam.

Ministry of Health. (1997b). **National Control Program HIV/AIDS/STD Surveillance of 1996**. Epidemiology Unit, National Aids Control Program. Report No. 11. Dar es Salaam: Government Printer.

Ministry of Health. (2006). National Guidelines for Malaria Diagnosis and Treatment. **Malaria Control Series 11**. Dar es Salaam: Government Printer.

Mkomozi Centre. (2005). **Demonstrating the potential of MEMKWA to respond to the learning needs of street children**. Accessed 22 May 2006 at <http://www.eldis.org/cf/search/disp/DocDisplay.cfm?Doc=DOC22206&Resource=f1educ>

Mujinja. (2000). Equity of access to public, private not-for-profit and private for-profit health facilities in two regions of Tanzania. Muhimbili University College of Health Science, Dar es Salaam. Unpublished.

Muli J. (2000). **Street children programme: Undugu Annual Report 1999**. Accessed 20 July 2004 at http://home01.wxs.nl-heinie/assets/annual_report_1999_STREE_CHILDREN_PROGRAMME.pdf

Murdock O P, Garbharran H, Edwards J M, Smith A M, Lutcmiah J and Mkhize M. (2003). Peer-led HIV/AIDS prevention for women in South African informal settlements. **Health Care for Women International**, 24: 502-512.

Mvungi A K. (2002). The state and status of street children in Tanzania today. A theoretical framework. In: Lugalla J P and Kibassa C G. (Eds) **Poverty, Aids and street children in East Africa**. Lewiston, New York: Mellen Press.

Mwagani A E, Ankomah A and Powell A R. (2001). Sex preference and contraceptive behaviour among men in Mbeya region, Tanzania: A rural-urban comparison of qualitative data. **Journal of Biosocial Science**, 30: 381-382.

Nsimba S. (2006). How sulfadoxine-pyrimethamine (SP) was perceived in some rural communities after phasing out chloroquine (CQ) as a first-line drug for uncomplicated malaria in Tanzania: Lessons to learn towards moving from monotherapy to fixed combination therapy. **Online Journal of Ethnobiology and Ethnomedicine**. Available at <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1379633&rendertype>

Nsungwa-Sabiiti J, Kallander K and Nsabagasani. (2004). Local fever illness classifications: Implications for home management of malaria strategies. **Tropical Medicine International Health**, 9: 1191-1199.

Ntarangwi M. (2003). **Report of ten years of Street Children Programme (1987-1997)**. Plan International. Embu – Meru Nakuru, Kenya: Media Document Supplies.

- Ntozi M P J. (1997). Effects of AIDS on children: The problem of orphans in Uganda. *Health Transition Review*, 7 (Supplement): 23-40. Accessed 20 June 2005 at <http://htc.anu.edu.au/pdfs/Ntozi3.pdf>
- Philemon, J.R.M and Kessy S.S.A. (2008). *Negotiating safe sex among women: The fight against HIV/AIDS in Tanzania*. Research Report 08.5, Dar es Salaam: REPOA.
- Population Reports. (2001). *Youth and HIV/AIDS: Can we avoid catastrophe*. Report No.26 (3). Accessed 20 April 2004 at <http://www.infoforhealth.org/pr/112/112print.shtml>
- Prohaska R T, Albrecht G, Levy A J, Sugrues N and Kim J. (1990). Determinants of self-perceived risk for AIDS. *Journal of Health and Social Behaviour*, 31: 384-394.
- Raffaelli M, Campos R, Merritt P A, Siqueira E, Antunes M C, Parker R, Greco D, Halsey N, Holt E, Lucio J, Kendall C, Ottoni Z, Andrade P, Rolf J, Ude W and de Zalduondo B. (1993). Sexual practices and attitudes of street youth in Belo Horizonte, Brazil. *Social Science and Medicine*, 37 (5): 661-670.
- Rajani R and Kudrati M. *The variety of sexual experience of street children in Mwanza and their implications for sex education/HIV prevention programs*. Kuleana Center for Children's Rights.
- Rajani R, Bangser M, Lund-Sorensen U and Leach V. (2001). Situation analysis of children in Tanzania. *Tanzania Online*. Accessed 30 July 2005 at <http://www.eldis.org/cf/search/disp/OrgDisplay.cfm?Org=5710&Rsource=f1>
- Rau B. (2002). *Combating child labour and HIV/AIDS in sub-Saharan Africa: A review of policies, programs, and projects in South Africa*.
- Redman J. (1987). AIDS and peer teaching. *Health Education Journal*, 46 (4): 150-151.
- Tanzania and Zambia to identify good practices, Paper No 1*. International Labour Organization, International Program on the Elimination of Child Labor (IPEC).
- Rotheram-Borus M, Becker J, Koopman C and Kaplan M. (1991). AIDS knowledge and beliefs, and sexual behaviour of sexuality delinquent and non-delinquent (runaway) adolescents. *Journal of Adolescence*, 24: 229-244.
- Ruto S. (1999). Why we go to the street: A baseline study on the phenomenon of Nairobi's street children. Unpublished report.
- Social Action Trust Fund. (2006). Annual program review 2005. Unpublished report.
- Save the Children. (2003). *The unbearable cost of illness: Poverty, ill-health and access to health care-evidence from Lindi Rural District, Tanzania*. Accessed 11 October 2006 at <http://www.savethechildren.org.uk/cocwi/Tanzania.pdf>
- Scanlon J T, Tomkins A, Lynch A M and Scanlon F. (1998). Street children in Latin America. *British Medical Journal*, 316: 1596-1600.
- Seligman M. (1992). *Helplessness*. Freeman: New York.
- Shuey A D, Babishangire B B, Omist S and Bagarukayo H. (1999). Increased sexual abstinence among in-school adolescents as a result of school health education in Soroti district, Uganda. *Health Education Research*, 4 (3): 411-419.
- Situmorang A. (2005). *Sex and sexually transmitted infections including HIV/AIDS among street youth in Indonesia*. A research proposal. Accessed 20 June 2005 at http://www.gfmer.ch/Medical_education_En/PGC_RH_2005/pdf/Street_youth_Indonesia_R.pdf

- Swart-Kruger J and Richter M L. (1997). AIDS-related knowledge, attitudes and behaviour among South African street youth: Reflections on power, sexuality and the autonomous self. ***Social Science and Medicine***, 45 (6): 957-966.
- Tanzania Commission for AIDS (TACAIDS). (2003). ***National Multi-sectoral Strategic Framework on HIV/AIDS 2003-2007***. Accessed 1 July 2004 at <http://www.tanzania.go.tz/pdf/tacaidsnmf.pdf>
- Tsikata. (1995). ***Effects of structural adjustment on women and the poor***. Accessed 16 December 2003 at <http://www.twinside.org.sg/souths/twn/women.htm>
- UNAIDS. (2006). ***Report on the global AIDS epidemics***. Accessed 21 September 2006 at <http://www.unaids.org/en/HIVdata/2006GlobalReport/default.asp>
- UNAIDS/UNICEF/WHO. (2002). ***United Republic of Tanzania: Epidemiological fact sheets on HIV/AIDS and sexually transmitted diseases***. Accessed 10 April 2004 at http://www.who.int/emc-hiv/fact_sheets/pdfs/Tanzania_EN.pdf
- UNICEF. (1989). ***Annual Report***. New York: United Nations.
- UNICEF. (1990). ***Women and children in Tanzania: A situational analysis***. Dar es Salaam: UNICEF.
- UNICEF. (2001). ***A study of street children in Zimbabwe***. Accessed 20 September 2005 at <http://www.unicef.org/reseval/evaldb/street%20children.pdf>
- UNICEF. (2003). ***Africa's orphaned generations***. Accessed 29 May 2003 at http://www.unicef.org/publications/index_16271.html
- United Republic of Tanzania. (2002). ***Population and Housing Census***. Accessed on 20 June 2008 at <http://www.tanzania.go.tz/census.htm>
- Volpi E. (2003). Street children promises and approaches. ***Children, Youth and Environment***, 13 (1). Accessed 14 July 2004 at <http://cye.colorado.edu/Vol113ArticleReprints/PromisigPractices.pdf>
- Women's Dignity Project. (2004). ***Poor people experiences of health services in Tanzania***. Accessed 14 May 2007 at http://www.tanzaniagateway.org/docs/poor_peoples_experiences_of_health_services_in_Tanzania.pdf
- World Health Organisation (WHO). (1993). ***A one-way street? Report on Phase 1 of the street children project: Part 2 of 9 - Programme on substance abuse***. Accessed 27 May 2004 at http://pangaea.org/street_children/world/who2.htm
- World Health Organization (WHO). (2000). ***Working with street children. Module 7, Teaching street children: A training package on substance abuse, sexual and reproductive health including HIV and STDs***. Accessed 12 July 2004 at http://whqlibdoc.who.int/hq/2000/WHO_MSD_00.14_Module7.pdf
- Yamamoto R. (1996). ***Street girls in Dar es Salaam: A situational analysis***. Dar es Salaam: Africa Education Fund International.

Appendices

Appendix 1: Findings from the Control Sample

To ensure the reliability of survey results, data were collected from a control sample of 40 boys who had previously worked and/or lived on the streets. Tables A1 and A2 present findings from the control sample, and the results are compared with survey data below.

Table A1: General Characteristics of the Control Sample

Variable	Respondents (N=40)	
	Frequency	Percentage
Age		
6-10	13	33
11-15	19	48
16-18	8	20
How long living / working in the street (years)		
Under 1 year	12	30.0
Over 1 year	28	70.0

Table A2: Key Findings from the Control Sample

Variable	Frequency	Percentage
Diseases that street children frequently suffer (N=40)		
Diarrhoea	14	35
Skin diseases	25	63
Chest diseases	13	33
Fever	22	55
Headache	17	43
Stomach ache	14	35
Accidents/Injury	21	53
Health-seeking behaviours / coping strategies (N=40)		
Going to hospital	11	28
Going to traditional healers	0	0
Sleep off the illness	8	20
Buying medicine	21	53
Reasons for buying medicine (n=21)		
Cheap	9	43
Saves time and nearby	7	33
Effective	5	24

Results from the control sample correspond to major findings among the survey population. Consistent with the survey population, the majority of respondents in the control sample had lived/worked on the streets for more than one year (control sample 70%; main survey, boys 61% and girls 54%).

Respondents in the control sample also frequently cited suffering various diseases/illnesses while on the streets including skin diseases, headaches, fever, accidents and injury, and diarrhoea. As found in the main survey, buying medicines from local pharmacies was the most common coping strategy for illness cited by the control respondents. Moreover, the reasons provided by respondents for this behaviour corresponded closely to the testimony provided by survey participants. In both control sample and survey, the two reasons most commonly cited for buying medicine were it was cheap and it saved time. By adopting this strategy, children could attend to their other immediate needs, such as looking for food and a place to sleep.

Appendix 2: Swahili Questionnaire

MBINU WANAZOTUMIA WATOTO WANAOSHII MITAANI KUJITIBU WANAPOKUA WAGONJWA

A. Habari za Msingi

Kata

Mtaa

Jinsia

Tarehe

1. Umri

- a. Miaka 6-10
- b. Miaka 11-15
- c. Miaka 16-18

2. Kwa kawaida huwa unalala wapi?

- a. Mtaani
- b. Nyumbani
- c. Mtaani na nyumbani
- d. Geto
- e. Sehemu nyingine (elezea)

3. Kama huwa unalala nyumbani, ni nani unayeishi naye?

- a. Wazazi wote wawili (baba na mama)
- b. Mama
- c. Baba
- d. Bibi
- e. Babu
- f. Wengine (elezea)

4. Je mama yako yuko hai?

- a. Ndio
- b. Hapana

5. Je baba yako yuko hai?

- a. Ndio
- b. Hapana

6. Kwa muda gain sasa umekua ukiishi/kufanya kazi mitaani?

- a. Kwa muda usiozidi mwaka mmoja
- b. Mwaka mmoja
- c. Zaidi ya mwaka mmoja

7. Ni kazi/ shughuli gain unayofanya mitaani?

- a. Kuuza chakula
- b. Kuuza magazeti
- c. Kuokota majalalani
- d. Kuomba omba
- e. Nyingine (elezea) _____

8. Kiwango cha elimu

- a. Hajasoma
- b. Hajamaliza elimu ya msingi
- c. Bado anasoma
- d. Amemaliza elimu ya msingi
- e. Elimu ya Sekondari

B. Magonjwa yanayowashambulia watoto waishio mitaani

9. Je umeshawahi kuumwa kwa kipindi cha miezi mitatu iliyopita?
- Ndio
 - Hapana

Nimagonjwa gain uliyomwa? (Usisome majibu)

Ugonjwa	Ndio	Hapana	Idadi ya siku
Kuharisha			
Magonjwa ya ngozi			
Kifua/kukohoa			
Malaria			
Homa			
Tumbo			
Kichwa			
Ajali/ vidonda			
Kuumwa na nyoka			
Mwingine (elezea)			

C. Mbinu za kujitibu

10. Kama ulishawahi kuumwa magonjwa yeyote kati ya hayo uliyotaja kwenye swali la 9, ulifanya nini? (Usisome majibu)
- Nilienda hospitali
 - Nililala
 - Nilienda kwa waganga wa kienyeji
 - Nilinunua dawa
 - Mbinu nyingine (elezea)_____
11. (Kulingana na jibu la swali la 10) Kwa nini ulichagua mbinu hiyo?
-
 -
 -
 -
12. (Kulingana na majibu ya swali la 10) kwa nini haukuchagua kwenda hospitali? hospitali?
-
 -
 -
 -
13. (Kulingana na majibu ya swali la 10) nani aliyekushauri kuchagua mbinu hiyo?
- Mimi mwenyewe
 - Rafiki yangu
 - Ndugu yangu
 - Mwingine (mtaje)_____

14. Kama ulienda hospitali, ni nani alikupeleka??
 - a. Mimi mwenyewe
 - b. Rafiki yangu
 - c. Mama
 - d. baba
 - e. Mwingine (mtaje)_____
 - d. Others (specify)

15. Kama ulikwenda hospitali, ulipata matatizo yeyote katika kupata matibabu?
 - a. Ndio
 - b. Hapana

16. Kama ndiyo (kulingana na swali la 15) ni matatizo gain uliyokutana nayo? (Uisome majibu)
 - a. Sikuwa na pesa za kutosha kununua dawa
 - b. Nilifukuzwa nisiingie hospitali
 - c. Watumishi wa hospitali walikua wakatili kwangu
 - d. Tatizo lingine (elezea)_____

17. Ili uweze kwenda hospitali na kupata matibabu, ungependa huduma za afya zeweje?
 - a.
 - b.
 - c.
 - d.

D. Maswali kuhusu Tabia zinazoweza kufanya kuambukizwa UKIMWI

18. Je unajua kondomu ni nini?
 - a. Ndio
 - b. Hapana

19. Umeshawahi kusikia kuhusu kondomu?
 - a. Ndio
 - b. Hapana

20. Umeshawahi kutumia kondomu?
 - a. Ndio
 - b. Hapana

21. Ni sababu gain zilizokufanya utumie kondomu?
 - a. Kuzuia mimba
 - b. Kuzuia kumpa mamba mpenzi wangu
 - c. Kuzuia kuambukiza magonjwa
 - d. Kuzuia kuambukizwa magonjwa
 - e. Sababu nyingine(elezea)_____

22. Umeshawahi kufanya ngono?
 - a. Ndio
 - b. Hapana

23. Je, ulitumia kondomu mara ya mwisho ulipofanya ngono?
 - a. Ndio
 - b. Hapana (Nenda swali namba 24)

24. Kwa nini hukutumia kondomu?
- Mpenzi wangu alikataa
 - Sikujua kuhusu kondomu
 - Kondomu inazuia kufurahia ngono
 - Niliona aibu kumwambia mpenzi wangu
 - Sababu nyingine (elezea)
25. Je unafanya kitu chochote kujikinga na UKIMWI?
- Ndio
 - Hapana
26. Kama ndio, unafanaya nini kujikinga na ugonjwa wa UKIMWI? (Usisome majibu)
- Kutumia kondomu
 - Kuepuka ngono
 - Kujitibu mapema magonjwa ya ngono
 - Sijui
 - Njia nyingine(elezea)_____

Appendix 3: English Questionnaire

Name of Research Assistant _____ Date _____

District _____ Place _____

COPING STRATEGIES USED BY STREET CHILDREN INCASE OF ILLNESS

Questionnaire

B. Social Demographic data

Ward Street

Sex.....Date.....

- Age
 - 6-10 years
 - 11-15 years
 - 16-18 years
- Where do you usually sleep (Probe but do not read out list)
 - In the street
 - At home
 - In the street and home
 - Ghetto (not at home not in the street)
 - Others (specify)_____
- If you sleep at home, with whom do you live with whom do you live with?
 - Both parents (mother and father)
 - Mother
 - Father
 - Grandmother/
 - Grandfather
 - Others (specify)_____
- Is your mother alive?
 - Yes
 - No
- Is your father alive?
 - Yes
 - No

6. For how long have you lived/worked in the street
 - a. Less than one year
 - b. One year
 - c. More than one year
 - d. Others (specify)_____
7. What type of work are you doing in the street?
 - a. Food vending
 - b. Selling magazine
 - c. Scavenging
 - d. Others (specify)_____
8. Level of education
 - a. No education
 - b. Have not completed primary school
 - c. Still in school
 - d. Completed primary education
 - e. Secondary education

B. Different Diseases experienced by street children

9. Have you ever been sick for the past six months?
 - a. Yes
 - b. No

What did you suffer from and for how long? (Probe but do not read out list)

Diseases	YES	NO	Number of Days
Diarrhoea			
Skin diseases			
Chest diseases			
Malaria			
Fever			
Stomach ache			
Head ache			
Accidents/injury			
Snake bite			
Others (specify)			

C. Coping strategies used and Health seeking behaviour

10. If you have ever suffered from any of the diseases in question 9, what did you do? (Probe but do not read the list)
 - a. Go to the hospital
 - b. Sleep
 - c. Go to the traditional healers
 - d. Buy medicine
 - e. (specify)

11. (According to the answer in question 10) why did you choose the mentioned alternative?
- a.
 - b.
 - c.
 - d.
12. (According to the answer in question 10) why didn't you go to the hospital?
- a.
 - b.
 - c.
 - d.
13. According to the answer in question 10) who advice you to choose the mentioned alternative?
- a. Myself
 - b. A friend
 - c. A relative
 - e. Others (specify)
14. If you went to the hospital, who took you to the hospital?
- a. Myself
 - b. My friend
 - c. My mother
 - d. My father
 - e. My relative (mention)_____
 - d. Others (specify)
15. If you went to the hospital, did you face any problems in getting treatment?
- a. Yes
 - b. No
16. If yes, (according to question 15) what problems did you face in getting treatment?
(Probe but do not read the list)
- a. I didn't have enough money to buy medicine
 - b. I was chased out of the hospital
 - c. hospital staffs were unfriendly
 - d. Other (specify)_____
17. In order for you to go to the hospital and access the treatment, how would like the health services to be?
- a.
 - b.
 - c.
 - d.

D. Behavioral questions

18. Have you ever heard about AIDS?
 - e. Yes
 - f. No
19. Have you ever heard about condom?
 - g. Yes
 - h. No
20. Have you ever used condom?
 - i. Yes
 - j. No
21. What was the main reason why you used condom? (unprompted)
 - k. To avoid pregnancy
 - l. To prevent making partner pregnant
 - m. To avoid giving diseases
 - n. To avoid getting diseases
 - o. Others (specify)_____
22. Have you ever had sex?
 - c. Yes
 - d. No
23. Did you use condom last time you had sex?
 - i. Yes
 - ii. No (Go to question 24)
24. Why didn't you use condom? (Unprompted)
 - iii Partner refusal
 - iv. I didn't know about condom
 - v. Condom limits feeling
 - vi. Too embarrassed to talk to the partner about condom
 - vii. Other reason (mention)
25. Are you doing anything to protect yourself from AIDS?
 - a) Yes
 - b) No
26. If yes, what are you doing to protect yourself against AIDS?
if (Probe but do not read out list)
 - a) Use condoms
 - b) Avoid sex
 - c) Early treatment of STDs
 - d) Don't know
 - e) Other (mention)_____

Appendix 4: Semi Structured Interview Guide

Topic: Coping strategies used by street children incase of illness

Good morning/afternoon / evening. Thank you so much for accepting to participate in this discussion. I would like to ask you some questions regarding issues of accessibility to health care, treatment and experiences in ill-health

I will appreciate if you could provide me with honest information and feel free to discuss your opinions. There is no right and wrong answer or opinion so be free to bring forward your thoughts, experiences and explanations in relation to this subject and I assure you that the information you give will remain confidential.

General Information:

1. What is your name? (Not necessary)
2. How old are you?
3. Sex _____
4. Are your parents alive?

Questions

A. Health seeking behavior/ Coping strategies

1. Where do you normally go to the hospital when you are sick?

Probe questions

- Why do you go to that place (according to response)
- Why don't you go to the hospital (if he/she go to the hospital)
- Who take you to the hospital/ other place where he/she seek treatment
- Who do you tell when you feel you are sick?
- Where do you get money for treatment?

B. Kind of diseases that affect street children most

2. What type of diseases or health problems you have experienced in the last two months?

Probe questions

- How many days didn't you work or play in the street as the result of that disease?
- What did you do?
- How did you know that you were sick?

3. For you to go to the hospital when you are sick, what would you like the health services to be?

C. Behavioral questions

4. What do you do to protect yourself from diseases?

Probe questions

- Have you ever heard about AIDS?
- Have you ever heard about condom?
- Have you ever used condom?
- How do you protect yourself from AIDS, STDs, Malaria, skin infections etc.
- Have you ever sex?

D. Sources of cash?

- What kind of work are you involving in the street?
- How much do you earn per day?


Thank you very much for your time. This is the end of the discussion.

Appendix 5: Participatory exercises guideline

Exercise 1: The spider diagram of street children experiences of health/diseases in their homes and while on the streets

- This exercise will involve two groups (one of street girls and the other of street boys)
- The street children will be given flip charts and marker pens and we will be told to draw a diagram of spider.
- The researcher will explain to street children that inside the body of the spider is the illness (the child will be told to write) and the outer circle is divided into two parts (home and streets). Each participant will write the diseases they have experienced in the past two months in the spider legs (see figure 1. below).

Figure 1: Spider diagram



- Using the health complaints mentioned, a table will be made and, by group consensus, they will rank the most frequent diseases attacking them and the places where they go for treatment or the coping mechanism they used during their sickness (see the table below). One child who knows how to write will be chosen to write down the scores agreed by the group.

Illness →								
Frequency								
Treatment sought/coping strategies								

Table to be completed by street children to show their illness and the action taken

- The discussions emerging from these exercises will be tape-recorded and the field notes will be taken.
- To researchers will be guiding the participants in this exercises.
- Each exercise is expected to take about 40 minutes.

Exercise 2: Mapping exercise to identify the places where street children go to seek treatment incase they are ill.

- This exercise will involve two groups (one boys and the other girls); each group will be consisted of 5 street children
- Street children will be given flip charts and maker pens and will be told to map the areas/places they go for treatment when they get sick
- The children will be told to explain what the map/drawing represents and the discussions that will emerge during the exercise will be tape-recorded and the field notes will be taken
- The exercise is expected to take about 30 minutes

Exercise 3: Mapping/drawing exercises to show the barriers the children are facing in accessing health services

- This exercise will involve two groups (one boys and the other girls); each group will be consisted of 5 street children
- Street children will be given flip charts and maker pens and will be told to map/draw the barriers they face in accessing the health services.
- The children will be told to explain what the map/drawing represents and the discussions that will emerge during the exercise will be tape-recorded and the field notes will be taken
- The exercise is expected to take about 30 minutes

Exercise 4: Mapping/Drawing exercise to show what the children do when they get Sick

- This exercise will involve two groups (one boys and the other girls); each group will be consisted of 5 street children
- Street children will be given flip charts and maker pens and will be told to map/draw the barriers they face in accessing the health services.
- The children will be told to explain what the map/drawing represents and the discussions that will emerge during the exercise will be tape-recorded and the field notes will be taken
- The exercise is expected to take about 30 minutes

Publications by REPOA

Books

"Researching Poverty in Tanzania: problems, policies and perspectives"

Edited by Idris Kikula, Jonas Kipokola, Issa Shivji, Joseph Semboja and Ben Tarimo

"Local Perspectives on Globalisation: The African Case"

Edited by Joseph Semboja, Juma Mwapachu and Eduard Jansen

"Poverty Alleviation in Tanzania: Recent Research Issues" Edited by M.S.D. Bagachwa

Research Reports

10/1 *"Coping Strategies Used by Street Children in the Event of Illness"*
Zena Amury and Aneth Komba

08.6 *"Assessing the Institutional Framework for Promoting the Growth of MSEs in Tanzania; The Case of Dar es Salaam"*
Raymond Mnenwa and Emmanuel Maliti

08.5 *"Negotiating Safe Sex among Young Women: the Fight against HIV/AIDS in Tanzania"*
John R.M. Philemon and Severine S.A. Kessy

08.4 *"Establishing Indicators for Urban Poverty-Environment Interaction in Tanzania: The Case of Bonde la Mpunga, Kinondoni, Dar es Salaam"*
Matern A.M. Victor, Albinus M.P. Makalle and Neema Ngware

08.3 *"Bamboo Trade and Poverty Alleviation in Ileje District, Tanzania"*
Milline Jethro Mbonile

08.2 *"The Role of Small Businesses in Poverty Alleviation: The Case of Dar es Salaam, Tanzania"*
Raymond Mnenwa and Emmanuel Maliti

08.1 *"Improving the Quality of Human Resources for Growth and Poverty Reduction: The Case of Primary Education in Tanzania"*
Amon V.Y. Mbelle

07.2 *"Financing Public Health Care: Insurance, User Fees or Taxes? Welfare Comparisons in Tanzania"*
Deograsias P. Mushi

07.1 *"Rice Production in the Maswa District, Tanzania and its Contribution to Poverty Alleviation"*
Jerry A. Ngailo, Abiud L. Kaswamila and Catherine J. Senkoro

06.3 *"The Contribution of Microfinance Institutions to Poverty Reduction in Tanzania"*
Severine S.A. Kessy and Fratern M Urio

06.2 *"The Role of Indigenous Knowledge in Combating Soil Infertility and Poverty in the Usambara Mountains, Tanzania"*
Juma M. Wickama and Stephen T. Mwihomeke

06.1 *"Assessing Market Distortions Affecting Poverty Reduction Efforts on Smallholder Tobacco Production in Tanzania"*
Dennis Rweyemamu and Monica Kimaro

05.1 *"Changes in the Upland Irrigation System and Implications for Rural Poverty Alleviation. A Case of the Ndiwa Irrigation System, West Usambara Mountains, Tanzania"*
Cosmas H. Sokoni and Tamilwai C. Shechambo

04.3 *"The Role of Traditional Irrigation Systems in Poverty Alleviation in Semi-Arid Areas: The Case of Chamazi in Lushoto District, Tanzania"*
Abiud L. Kaswamila and Baker M. Masuruli

- 04.2 *"Assessing the Relative Poverty of Clients and Non-clients of Non-bank Micro-finance Institutions. The case of the Dar es Salaam and Coast Regions"*
Hugh K. Fraser and Vivian Kazi
- 04.1 *"The Use of Sustainable Irrigation for Poverty Alleviation in Tanzania. The Case of Smallholder Irrigation Schemes in Igurusi, Mbarali District"*
Shadrack Mwakalila and Christine Noe
- 03.7 *"Poverty and Environment: Impact analysis of Sustainable Dar es Salaam Project on "Sustainable Livelihoods" of Urban Poor"*
M.A.M. Victor and A.M.P. Makalle
- 03.6 *"Access to Formal and Quasi-Formal Credit by Smallholder Farmers and Artisanal Fishermen: A Case of Zanzibar"*
Khalid Mohamed
- 03.5 *"Poverty and Changing Livelihoods of Migrant Maasai Pastoralists in Morogoro and Kilosa Districts"*
C. Mung'ong'o and D. Mwamfupe
- 03.4 *"The Role of Tourism in Poverty Alleviation in Tanzania"*
Nathanael Luvanga and Joseph Shitundu
- 03.3 *"Natural Resources Use Patterns and Poverty Alleviation Strategies in the Highlands and Lowlands of Karatu and Monduli Districts – A Study on Linkages and Environmental Implications"*
Pius Zebbe Yanda and Ndalaha Faustine Madulu
- 03.2 *"Shortcomings of Linkages Between Environmental Conservation and Poverty Alleviation in Tanzania"*
Idris S. Kikula, E.Z. Mnzava and Claude Mung'ong'o
- 03.1 *"School Enrolment, Performance, Gender and Poverty (Access to Education) in Mainland Tanzania"*
A.V.Y. Mbelle and J. Katabaro
- 02.3 *"Poverty and Deforestation around the Gazetted Forests of the Coastal Belt of Tanzania"*
Godius Kahyarara, Wilfred Mbowe and Omari Kimweri
- 02.2 *"The Role of Privatisation in Providing the Urban Poor Access to Social Services: the Case of Solid Waste Collection Services in Dar es Salaam"* Suma Kaare
- 02.1 *"Economic Policy and Rural Poverty in Tanzania: A Survey of Three Regions"*
Longinus Rutasitara
- 01.5 *"Demographic Factors, Household Composition, Employment and Household Welfare"*
S.T. Mwisomba and B.H.R. Kiilu
- 01.4 *"Assessment of Village Level Sugar Processing Technology in Tanzania"*
A.S. Chungu, C.Z.M. Kimambo and T.A.L. Bali
- 01.3 *"Poverty and Family Size Patterns: Comparison Across African Countries"*
C. Lwechungura Kamuzora
- 01.2 *"The Role of Traditional Irrigation Systems (Vinyungu) in Alleviating Poverty in Iringa Rural District"*
Tenge Mkavidanda and Abiud Kaswamila
- 01.1 *"Improving Farm Management Skills for Poverty Alleviation: The Case of Njombe District"*
Aida Isinika and Ntengua Mdoe
- 00.5 *"Conservation and Poverty: The Case of Amani Nature Reserve"*
George Jambiya and Hussein Sosovele
- 00.4 *"Poverty and Family Size in Tanzania: Multiple Responses to Population Pressure?"*
C.L. Kamuzora and W. Mkanta

- 00.3 “*Survival and Accumulation Strategies at the Rural-Urban Interface: A Study of Ifakara Town, Tanzania*”
Anthony Chamwali
- 00.2 “*Poverty, Environment and Livelihood along the Gradients of the Usambaras on Tanzania*”
Adolfo Mascarenhas
- 00.1 “*Foreign Aid, Grassroots Participation and Poverty Alleviation in Tanzania: The HESAWA Fiasco*” S. Rugumamu
- 99.1 “*Credit Schemes and Women’s Empowerment for Poverty Alleviation: The Case of Tanga Region, Tanzania*”
I.A.M. Makombe, E.I. Temba and A.R.M. Kihombo
- 98.5 “*Youth Migration and Poverty Alleviation: A Case Study of Petty Traders (Wamachinga) in Dar es Salaam*”
A.J. Liviga and R.D.K Mekacha
- 98.4 “*Labour Constraints, Population Dynamics and the AIDS Epidemic: The Case of Rural Bukoba District, Tanzania*”
C.L. Kamuzora and S. Gwalema
- 98.3 “*The Use of Labour-Intensive Irrigation Technologies in Alleviating Poverty in Majengo, Mbeya Rural District*”
J. Shitundu and N. Luvanga
- 98.2 “*Poverty and Diffusion of Technological Innovations to Rural Women: The Role of Entrepreneurship*”
B.D. Diyamett, R.S. Mabala and R. Mandara
- 98.1 “*The Role of Informal and Semi-Formal Finance in Poverty Alleviation in Tanzania: Results of a Field Study in Two Regions*”
A.K. Kashuliza, J.P. Hella, F.T. Magayane and Z.S.K. Mvena
- 97.3 “*Educational Background, Training and Their Influence on Female-Operated Informal Sector Enterprises*”
J. O’Riordan. F. Swai and A. Rugumyamheto
- 97.2 “*The Impact of Technology on Poverty Alleviation: The Case of Artisanal Mining in Tanzania*”
B W. Mutagwaba, R. Mwaipopo Ako and A. Mlaki
- 97.1 “*Poverty and the Environment: The Case of Informal Sandmining, Quarrying and Lime-Making Activities in Dar es Salaam, Tanzania*”
George Jambiya, Kassim Kulindwa and Hussein Sosovele

Special Papers

- 09.31 “*Institutional Analysis of Nutrition in Tanzania*”
Valerie Leach and Blandina Kilama
- 09.30 “*Influencing Policy for Children in Tanzania: Lessons from Education, Legislation and Social Protection*”
Masuma Mamdani, Rakesh Rajani and Valerie Leach with Zubeida Tumbo-Masabo and Francis Omondi
- 09.29 “*Maybe We Should Pay Tax After All? Citizens’ Views of Taxation in Tanzania*”
Odd-Helge Fjeldstad, Lucas Katera and Erasto Ngalewa
- 09.28 “*Outsourcing Revenue Collection to Private Agents: Experiences from Local Authorities in Tanzania*”
Odd-Helge Fjeldstad, Lucas Katera and Erasto Ngalewa
- 08.27 “*The Growth – Poverty Nexus in Tanzania: From a Developmental Perspective*”
Marc Wuyts

08.26	<i>"Local Autonomy and Citizen Participation In Tanzania - From a Local Government Reform Perspective."</i> Amon Chaligha	17	<i>"When Bottom-Up Meets Top-Down: The Limits of Local Participation in Local Government Planning in Tanzania"</i> Brian Cooksey and Idris Kikula
07.25	<i>"Children and Vulnerability In Tanzania: A Brief Synthesis"</i> Valerie Leach	16	<i>"Local Government Finances and Financial Management in Tanzania: Observations from Six Councils 2002 – 2003"</i> Odd-Helge Fjeldstad, Florida Henjewe, Geoffrey Mwambe, Erasto Ngalewa and Knut Nygaard
07.24	<i>"Common Mistakes and Problems in Research Proposal Writing: An Assessment of Proposals for Research Grants Submitted to Research on Poverty Alleviation REPOA (Tanzania)."</i> Idris S. Kikula and Martha A. S. Qorro	15	<i>"Poverty Research in Tanzania: Guidelines for Preparing Research Proposals"</i> Brian Cooksey and Servacius Likwelle
07.23	<i>"Guidelines on Preparing Concept Notes and Proposals for Research on Pro-Poor Growth and Poverty in Tanzania"</i>	14	<i>"Guidelines for Monitoring and Evaluation of REPOA Activities"</i> A. Chungu and S. Muller-Maige
07.22	<i>"Local Governance in Tanzania: Observations From Six Councils 2002-2003"</i> Amon Chaligha, Florida Henjewe, Ambrose Kessy and Geoffrey Mwambe	13	<i>"Capacity Building for Research"</i> M.S.D. Bagachwa
07.21	<i>"Tanzanian Non-Governmental Organisations – Their Perceptions of Their Relationship with the Government of Tanzania and Donors, and Their Role and Impact on Poverty Reduction and Development"</i>	12	<i>"Some Practical Research Guidelines"</i> Brian Cooksey and Alfred Lokuji
06.20	<i>"Service Delivery in Tanzania: Findings from Six Councils 2002-2003"</i> Einar Braathen and Geoffrey Mwambe	11	<i>"A Bibliography on Poverty in Tanzania"</i> B. Mutagwaba
06.19	<i>"Developing Social Protection in Tanzania Within a Context of Generalised Insecurity"</i> Marc Wuyts	10	<i>"An Inventory of Potential Researchers and Institutions of Relevance to Research on Poverty in Tanzania"</i> A.F. Lwaitama
06.18	<i>"To Pay or Not to Pay? Citizens' Views on Taxation by Local Authorities in Tanzania"</i> Odd-Helge Fjeldstad	9	<i>"Guidelines for Preparing and Assessing REPOA Research Proposals"</i> REPOA Secretariat and Brian Cooksey
		8	<i>"Social and Cultural Factors Influencing Poverty in Tanzania"</i> C.K. Omari
		7	<i>"Gender and Poverty Alleviation in Tanzania: Issues from and for Research"</i> Patricia Mbughuni
		6	<i>"The Use of Technology in Alleviating Poverty in Tanzania"</i> A.S. Chungu and G.R.R. Mandara

- | | | | |
|---|--|----------|---|
| 5 | <i>“Environmental Issues and Poverty Alleviation in Tanzania”</i>
Adolfo Mascarenhas | Brief 12 | Changes in Citizens’ Perceptions of the Local Taxation System in Tanzania |
| 4 | <i>“Implications of Public Policies on Poverty and Poverty Alleviation: The Case of Tanzania”</i>
Fidelis Mtatifikolo | Brief 11 | Citizens Demand Tougher Action on Corruption in Tanzania |
| 3 | <i>“Who’s Poor in Tanzania? A Review of Recent Poverty Research”</i>
Brian Cooksey | Brief 10 | Outsourcing Revenue Collection: Experiences from Local Government Authorities in Tanzania |
| 2 | <i>“Poverty Assessment in Tanzania: Theoretical, Conceptual and Methodological Issues”</i>
J. Semboja | Brief 9 | Children and Vulnerability in Tanzania: A Brief Overview |
| 1 | <i>“Changing Perceptions of Poverty and the Emerging Research Issues”</i>
M.S.D. Bagachwa | Brief 8 | Mawazo ya AZISE za Tanzania Kuhusu Uhusiano Wao na Wafadhili |
| | | Brief 7 | Mawazo ya AZISE za Tanzania Kuhusu Uhusiano Wao na Serikali |
| | | Brief 6 | Local Government Reform in Tanzania 2002 - 2005: Summary of Research Findings on Governance, Finance and Service Delivery |

Project Briefs

- | | | | |
|----------|---|---------|---|
| Brief 18 | Planning in Local Government Authorities in Tanzania: Bottom-up Meets Top-down | Brief 5 | Children Participating in Research |
| Brief 17 | The Investment Climate in Tanzania: Views of Business Executives | Brief 4 | Changes in Household Non-Income Welfare Indicators - Can poverty mapping be used to predict a change in per capita consumption over time? |
| Brief 16 | Assessing the Institutional Framework for Promoting the Growth of Micro and Small Enterprises (MSEs) in Tanzania: The Case of Dar es Salaam | Brief 3 | Participatory Approaches to Local Government Planning in Tanzania, the Limits to Local Participation |
| Brief 15 | Preventing Malnutrition in Tanzania: A Focused Strategy to Improve Nutrition in Young Children | Brief 2 | Improving Transparency of Financial Affairs at the Local Government Level in Tanzania |
| Brief 14 | Influencing Policy for Children in Tanzania: Lessons from Education, Legislation and Social Protection | Brief 1 | Governance Indicators on the Tanzania Governance Noticeboard Website |
| Brief 13 | Disparities Exist in Citizens’ Perceptions of Service Delivery by Local Government Authorities in Tanzania | TGN1 | What is the Tanzania Governance Noticeboard? |
| | | LGR 12 | Trust in Public Finance: Citizens’ Views on taxation by Local Authorities in Tanzania |

- | | | | |
|--------|--|-------|--|
| LGR 11 | Domestic Water Supply: The Need for a Big Push | LGR 5 | Fees at the dispensary level: Is universal access being compromised? |
| LGR10 | Is the community health fund better than user fees for financing public health care? | LGR 4 | TASAF – a support or an obstacle to local government reform |
| LGR 9 | Are fees the major barrier to accessing public health care? | LGR 3 | Councillors and community leaders – partnership or conflict of interest? Lessons from the Sustainable Mwanza Project |
| LGR 8 | Primary education since the introduction of the Primary Education Development Plan | LGR 2 | New challenges for local government revenue enhancement |
| LGR 7 | Citizens' access to information on local government finances | LGR 1 | About the Local Government Reform Project |
| LGR 6 | Low awareness amongst citizens of local government reforms | | |